

FORM PTO-1390
(REV. 1-98)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER
97 AE CNR IMMTRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/403627

INTERNATIONAL APPLICATION NO.
PCT/FR98/00923INTERNATIONAL FILING DATE
07 MAY 1998PRIORITY DATE CLAIMED
07 MAY 1997TITLE OF INVENTION
PEPTIDE ANALOGUES AND USES THEREOF, IN PARTICULAR IN PHARMACEUTICAL AND DIAGNOSTIC COMPOSITIONSAPPLICANT(S) FOR DO/EO/US Jean-Gerard GUILLET, Gilles GUICHARD, Maria OSTANKOVITCH,
Francine CONNAN, Anne QUESNEL, Jeannine CHOPPIN, Jean-Paul BRIAND and Sylviane MULLER

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: Patent Application Data Entry Format Sheet

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)

09/403627

INTERNATIONAL APPLICATION NO. PCT/FR98/00923

ATTORNEY'S DOCKET NUMBER
97 AE CNR IMM17. ☒ The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):

Neither international preliminary examination fee (37 CFR 1.482)
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO
and International Search Report not prepared by the EPO or JPO \$ 970.00International preliminary examination fee (37 CFR 1.482) not paid to
USPTO but International Search Report prepared by the EPO or JPO \$840.00International preliminary examination fee (37 CFR 1.482) not paid to USPTO
but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$760.00International preliminary examination fee (37 CFR 1.482) paid to USPTO
but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$670.00International preliminary examination fee (37 CFR 1.482) paid to USPTO
and all claims satisfied provisions of PCT Article 33(1)-(4) \$96.00

ENTER APPROPRIATE BASIC FEE AMOUNT =

CALCULATIONS PTO USE ONLY

\$ 840

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☒ 30
months from the earliest claimed priority date (37 CFR 1.492(e)).

\$ 130

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$
Total claims	20 - 20 =	0	x \$18.00	\$
Independent claims	4 - 3 =	1	x \$78.00	\$ 78
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$260.00	\$

TOTAL OF ABOVE CALCULATIONS =

\$ 1,048

Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement
must also be filed (Note 37 CFR 1.9, 1.27, 1.28).

+

SUBTOTAL =

\$ 1,048

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(f)).

\$

TOTAL NATIONAL FEE =

\$ 1,048

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +

\$

TOTAL FEES ENCLOSED =

\$ 1,048

Amount to be
refunded:

\$


charged:

\$

a. ☒ A check in the amount of \$ 1,048. to cover the above fees is enclosed.b. ☐ Please charge my Deposit Account No. in the amount of \$ to cover the above fees.
A duplicate copy of this sheet is enclosed.c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required by
37 CFR 1.16 and 1.17, or credit any overpayment to Deposit Account No. 25-0120. A duplicate
copy of this sheet is enclosed.NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR
1.137 (a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

November 5, 1999

Young & Thompson
745 South 23rd Street
2nd Floor
Arlington, VA 22202
(703) 521-2297


SIGNATURE

Andrew J. Patch
NAME32,925
REGISTRATION NUMBER

UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of
Jean-Gérard GUILLET et al.

Serial No. (unknown)

Filed herewith

PEPTIDE ANALOGUES AND USES THEREOF,
IN PARTICULAR IN PHARMACEUTICAL AND
DIAGNOSTIC COMPOSITIONS

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents

Washington, D.C. 20231

Sir:

Prior to the first Official Action and calculation
of the filing fee, please amend the above-identified applica-
tion as follows:

IN THE CLAIMS:

Claim 3, line 1, cancel "or 2".

Claim 4, line 1, change "one of Claims 1 to 3," to
--Claim 1,--.

Claim 5, line 1, change "one of Claims 1 to 4," to
--Claim 1,--.

Claim 6, line 1, change "one of Claims 1 to 5," to
--Claim 1,--.

Claim 10, line 1, change "one of Claims 1 to 9," to
--Claim 1,--.

Claim 11, line 2, change "one of Claims 1 to 9," to
--Claim 1,--.

Claim 12, line 2, change "one of Claims 1 to 9," to
--Claim 1,--.

Jean-Gérard GUILLET et al.

Claim 13, line 6, change "one of Claims 1 to 9," to
--Claim 1,--.

Claim 14, line 1, change "one of Claims 1 to 9," to
--Claim 1,--.

Claim 15, line 8, change "one of Claims 1 to 9," to
--Claim 1,--.


Claim 16, line 1, cancel "the";
line 2, cancel "according to Claim 15";
line 3, change "one of Claims 1 to 9;" to
--Claim 1;--;

line 6, change "the" to --a--; cancel
"according to Claim 14".

Claim 19, line 1, change "one of Claims 1 to 9," to
--Claim 1,--.

Respectfully submitted,

YOUNG & THOMPSON

By 
Andrew J. Patch
Attorney for Applicants
Registration No. 32,925
745 South 23rd Street
Arlington, VA 22202
Telephone: 703/521-2297

November 5, 1999

PEPTIDE ANALOGUES AND USES THEREOF, IN PARTICULAR IN PHARMACEUTICAL AND DIAGNOSTIC COMPOSITIONS

5 The present invention relates to peptide analogues and to their uses, mainly in preparing pharmaceutical compositions, in particular vaccines, and for *in vitro* or *in vivo* diagnosis of various pathologies.

10 The development of neuropeptides, peptide hormones and peptide-based antibiotics, or peptide-based synthetic vaccines, is greatly disrupted by the high sensitivity of peptides towards proteolysis, which limits, *inter alia*, their oral and parenteral administration.

15 Pseudopeptides are a class of molecules which are particularly advantageous in the design of enzyme inhibitors. Cleavage with peptidases is involved in a variety of biological processes, since it makes it possible to generate active peptide or protein fragments from inactive precursors. The idea that an isosteric bond could be considered as a mimic of the tetrahedral intermediate formed during passage through the transition state has opened the way to inhibitors of pseudopeptide type. At the start of the 1980s this approach was used successfully in the development of powerful inhibitors of angiotensin conversion enzyme (ACE), of renin (Szelke *et al.*, 1982 ; Boger *et al.*, 1983) and of the aspartic protease of HIV (Huff, 1991). More recently, several selective pseudopeptide inhibitors of Ras farnesyl transferase have been proposed. These molecule are of potential therapeutic value in the treatment of cancers involving the mutated Ras oncogene (Buss & Marsters, 1995).

25 Pseudopeptidiques analogues of peptide hormones have also been the subject of many studies, with the prime objective of stabilizing the starting peptide and conserving the activity. This strategy has effectively allowed the discovery of powerful agonists (Chorev *et al.*, 1979 ; Nagain *et al.*, 1988 ; Doulut *et al.*, 1992), but also of partial agonists and even antagonists of the hormone studied (Martinez *et al.*, 1985 ; Coy *et al.*, 1989 ; Leban *et al.*, 1993 ; Cai *et al.*, 1994). Analogues which are more selective at the receptor level have also been obtained in certain cases (Mendre, 1988 ; Nagain *et al.*, 1988 ; Schiller *et al.*, 1993). Moreover, the pseudopeptide bond makes it possible to explore the conformational characteristics of the ligand at the modified site and to evaluate the contribution of the amide bond and the inter- or intramolecular hydrogen bonding in the interaction with the receptor or receptors.

However, the modifications made hitherto on peptides for immunological use are limited essentially to the addition of sugars, to that of longer or shorter amino acids to allow the emulsification of the resulting lipopeptides, to cyclization by formation of disulphide or covalent bridges, as well as to the addition of molecules such as biotin, in order to increase, for example, the sensitivity of solid-phase immunochemical tests.

Several authors have asserted that pseudopeptides probably have very little or no immunogenicity, since they could not be transformed and presented to the molecules of the major histocompatibility complex (MHC) in order to be recognized by the helper T cells or by the cytotoxic T lymphocytes. In this respect, Dintzis *et al.* have described that the L enantiomer of rubredoxin induces a strong immune response by production of immunoglobulins of isotype G (IgG), whereas the corresponding protein consisting of amino acids all of D configuration does not induce an immune response.

It is in this context that the immunoretroids described below, as well as the antibodies directed against these immunoretroids, have been described in International Patent Application WO 95/24916 as being useful in vaccinating against pathologies in which peptides from which they are derived (parent peptides) were involved, as well as for the diagnosis of these pathologies.

The authors of that International Patent Application, who are also the authors of the present application, had, in point of fact, demonstrated for the first time that the modification by replacing a peptide bond -CO-NH- in the peptide chain of a peptide, with a bond -NH-CO-, and, where appropriate, by the replacement of an amino acid of L configuration with an amino acid of D configuration, makes it possible to obtain peptide analogues (retro or retro-inverso, respectively) which can be used in the treatment of pathologies involving the immune response to humoral or cell mediation.

The abovementioned immunoretroids are compounds of the peptide type, i.e. compounds consisting of a chain of protein-generating amino acids, in which at least one of the -CO-NH- bonds, and advantageously all of the -CO-NH- bonds, in the peptide chain of the corresponding parent peptide (not comprising any -NH-CO- bonds in its peptide chain) is (are) replaced with one (or more) -NH-CO- bonds, the chirality of each aminoacyl residue, whether or not it is involved in one or more of the abovementioned -NH-CO bonds, being either conserved (retro peptide) or inverted (retro-inverso peptide) relative to the corresponding aminoacyl residues constituting the said parent peptide.

The expression "retro-inverso peptide" should be understood as meaning any peptide or peptide analogue corresponding to the definition of immunoretroids given above, the said peptide consisting more particularly of a peptide chain in which at

least one of the residues is linked to at least one neighbouring residue via an -NH-CO- bond, on the one hand, and has the opposite chirality to that of this same aminoacyl residue in the peptide chain of the parent peptide, on the other hand.

The expression "retro peptide" should be understood as meaning any peptide corresponding to the definition of immunoretroids given above, the said peptide consisting more particularly of a peptide chain in which at least one of the residues is linked to at least one neighbouring residue via an -NH-CO- bond, the chirality of all of the aminoacyl residues involved in at least one -NH-CO- bond being conserved relative to the corresponding residue in the peptide chain of the parent peptide.

It goes without saying that the -CO-NH- et -NH-CO- bonds must be taken into account in the text hereinabove and hereinbelow, in the sense of the parent peptide chain going from the amino-terminal (N-terminal) end to the carboxy-terminal (C-terminal) end.

Given the general context of uncertainty regarding the efficacy of the peptide analogues recalled above, the authors of this International patent application WO 95/24916 considered that this inversion of the direction of the amide bond between protein-generating amino acids, and/or the modification of the configuration of the protein-generating amino acids, represented the only possible modifications of the parent peptides in order to obtain peptide analogues with a longer half-life, and biological properties, in particular immunological properties, which are comparable with, or even higher than, those of the abovementioned parent peptides.

The present invention derives from the discovery made by the authors of the present patent application that, contrary to what they believed they had correctly deduced in the context of their previous invention on account of the general context recalled above, these modifications of retro or retro-inverso type of the peptide chain of the parent peptides do not represent the only possible modifications for obtaining peptide analogues of interest, at least not in the context of the diagnosis and/or treatment of pathologies involving the immune response to cell mediation.

The reason for this is that the authors of the present patent application have demonstrated that the modification of natural peptides involved in pathologies in which immunity to cell mediation is involved, and more particularly the cytotoxic T lymphocytes, by replacing peptide bonds -CO-NH- in the peptide chain of the said natural peptides with bonds other than the -NH-CO- bonds, and/or by replacing the protein-generating amino acids in the natural peptide with non-protein-generating amino acids, makes it possible to obtain peptide analogues of the said natural peptides which can be used with the abovementioned advantages in the context of

the treatment or diagnosis of the said pathologies.

The expression "protein-generating amino acid" means, in the text hereinabove and hereinbelow, any amino acid which is involved in the construction of a natural peptide or protein.

5 The expression "non-protein-generating amino acid" means, in contrast with the above definition, any amino acid which is not involved in the construction of a natural peptide or protein. The expression "non-protein-generating amino acid" more particularly refers to any amino acid in which the carbon bearing the side chain R, i.e. the group -CHR-, located between -CO- and -NH- in the natural peptide chain, is
10 replaced with a unit which is not involved in the construction of a natural peptide or protein.

The object of the present invention is to provide pharmaceutical compositions, and more particularly vaccines, comprising peptide analogues which have a markedly longer half-life than that of natural peptides or natural proteins, or than the
15 half-life of synthetic peptides which may or may not be derived from these natural proteins (these natural proteins, or peptides which may or may not be derived from these natural proteins, also being referred to in the text hereinbelow by the expression "parent peptides or proteins"), of which they are analogues, while at the same time having biological activity, and more particularly immunological activity, which is comparable to, or even higher than, that of the abovementioned parent
20 peptides or proteins, or alternatively an activity which is different to or opposite that of the said parent peptides or proteins

Another object of the present invention is to provide methods for the *in vivo* diagnosis or alternatively for the *in vivo* evaluation of an individual's immune response capacity in the context of pathologies in which natural peptides or natural
25 proteins (which may be exogenous or endogenous) are liable to be involved by binding to the molecules of the MHC, the peptide-MCH complexes thus obtained [lacuna] to the receptors located on the T cells and recognizing the said natural peptides or natural proteins in their form complexed with the said molecules of the
30 MHC.

A further object of the invention is to provide compositions and kits for carrying out the abovementioned methods of *in vivo* diagnosis or evaluation of the immune response.

35 A further object of the present invention is to provide methods for the *in vitro* diagnosis of pathologies associated with the presence of endogenous or exogenous proteins in an individual's body, these methods being carried out with the aid of the peptide analogues as defined above, or with the aid of antibodies directed against the complexes between these antibodies and the molecules of the MHC, and having the advantage of giving higher-quality performance than the diagnostic methods

currently carried out with the aid of the parent peptides or proteins, or with the aid of antibodies directed against these parent peptides or proteins.

A further object of the present invention is to provide new kits for carrying out such methods of *in vitro* diagnosis.

5 The present invention relates to the use of peptide analogues of parent peptides, these parent peptides being derived, where applicable, from exogenous or endogenous proteins, the said parent peptides interacting with molecules of the MHC in the context of pathologies involving an immune response to cell mediation, in man or animals, the said analogues being characterized in that they correspond to
10 the said parent peptides in which:

- at least one peptide bond -CO-NH- of the peptide chain is modified, with the exception of modifications of the retro or retro-inverso type, or

- at least one amino acid of the peptide chain is substituted with a non-protein-generating amino acid, or

- 15 - at least one peptide bond -CO-NH- of the peptide chain is modified and at least one amino acid of the said peptide chain is substituted with a non-protein-generating amino acid,

for the preparation of a medicinal product intended for preventing or treating the abovementioned pathologies.

20 More particularly, the invention relates to the abovementioned use of peptide analogues derived from parent peptides which interact with molecules of the MHC of category I, in the context of pathologies involving cytotoxic T lymphocytes.

Advantageously, the peptide analogues used in the context of the present invention, and more particularly those derived from parent peptides which interact
25 with molecules of the MHC of category I, correspond to the said parent peptides in which at least one peptide bond -CO-NH- of the peptide chain is modified, with the exception of modifications of the retro or retro-inverso type.

As has already been seen previously, the abovementioned expression "parent peptide" should be understood as meaning,

- 30 - either a peptide which exists per se in the natural state, in particular in a microorganism or in a higher organism (in particular in the human body),

- or any peptide of immunological interest obtained by peptide synthesis, from protein-generating amino acids,

- 35 - or a peptide derived from a protein as exists in the natural state in the abovementioned organisms, in particular by fragmentation of the said protein (in particular with the aid of suitable proteinases, followed by purification of the peptide under consideration),

or by peptide synthesis (according to the methods conventionally used in this field),

- or a peptide derived from a protein as exists in the natural state, but whose immunological activity has been modified, conserved or optimized by replacing certain amino acids of the natural sequence with protein-generating amino acids, for example after screening a library of peptide analogues obtained by peptide synthesis.

The invention also relates to the peptide analogues of parent peptides, these parent peptides being derived, where applicable, from exogenous or endogenous proteins, it being possible for the said parent peptides to interact with molecules of the MHC in the context of pathologies in man or animals, the said analogues being characterized in that they correspond to the said parent peptides in which :

- at least one peptide bond -CO-NH- of the peptide chain is modified, with the exception of modifications of the retro or retro-inverso type, or

- at least one amino acid of the peptide chain is substituted with a non-protein-generating amino acid, or


- at least one peptide bond -CO-NH- of the peptide chain is modified and at least one amino acid of the peptide chain is substituted with a non-protein-generating [lacuna] acid.

Advantageously, the abovementioned peptide analogues of the invention are characterized in that the number of protein-generating or non-protein-generating aminoacyl residues, linked via a modified or unmodified bond, is between about 5 and about 20, preferably between 8 and 12 for the peptide analogues which bind to the molecules of the MHC of category I, and preferably between 8 and 16 for the peptide analogues which bind to the molecules of the MHC of category II.

More particularly, the invention relates to the peptide analogues as described above, characterized in that at least one of the peptide bonds -CO-NH- in the peptide chain of the parent peptide is replaced with a bond other than the -CO-NH- bond, the said other bond being chosen in particular from the following :

- | | |
|------------------------------------|-----------------------|
| $\text{-CH}_2\text{-NH-}$ | (methyleamino) ; |
| $\text{-CH}_2\text{-CH}_2\text{-}$ | (carba) ; |
| $\text{-CO-CH}_2\text{-}$ | (ketomethylene) ; |
| $\text{-CH}_2\text{-O-}$ | (methylenoxy) ; |
| $\text{-CHOH-CH}_2\text{-}$ | (hydroxyethylene) ; |
| -CHOH-CHOH- | (dihydroxyethylene) ; |
| -CH=CH- | (E or Z olefin) ; |
| -CHCN-NH- | (cyanomethyleamino) ; |
| $\text{-S-CH}_2\text{-}$ | (thiomethylene) ; |

-CH₂-S- (methylenethio) ;
 -CS-NH- (thioamide) ;
 -PO₂-NH- (phosphonamide) ;
 -CHOH- (hydroxymethylene) ;
 -NH-CO-NH- (urea) ;
 -CH₂-CH₂- (oxirane) ;



(tetrazole) ;

-CH₂-CO-NH- (β-homologation) ;
 -CHOH-CH₂-NH- (hydroxyethyleneamino) ;
 -CO-NH-NH- (hydrazino).

The invention also relates to the peptide analogues as described above, and characterized in that at least one of the peptide bonds -CO-NH- in the peptide chain of the parent peptide is replaced with a bond of the retro or retro-inverso type as defined above, in the case where at least one of the amino acids of the said peptide analogue is a non-protein-generating amino acid.

Peptide analogues which are preferred in the context of the present invention are characterized in that at least one of the peptide bonds -CO-NH- in the peptide chain or parent peptide is replaced with a methyleneamino bond or a bond of the β - homologation, or carba, or ketomethylene, or cyanomethyleneamino, or hydroxyethyleneamino type.

More particularly, the invention relates to the peptide analogues of parent peptides involved in melanoma, in particular the peptide MART1 27-35, the said parent peptide comprising, where appropriate, one or more mutations, such as the Leu²⁸-mutated parent peptide MART1 27-35, the said peptide analogues corresponding to the said parents peptides in which at least one of the peptide bonds -CO-NH- is modified, with the exception of modifications of the retro or retro-inverso type.

In this respect, the invention relates more particularly to the peptide analogues of the parent peptide Mart-1 27-35 of melanoma, comprising a methyleneamino bond and corresponding to the following formulae :

Sequences

5		P1	P2	P3	P4	P5	P6	P7	P8	P9
	MART1 27-35	H-A-	A-	G-	I-	G-	I-	L-	T-	V-OH
	Ψ (1-2)	H-A	Ψ(CH ₂ NH)A-	G-	I-	G-	I-	L-	T-	V-OH
	Ψ (2-3)	H-A-	A	Ψ(CH ₂ NH)G-	I-	G-	I-	L-	T-	V-OH
10	Ψ (3-4)	H-A-	A	G	Ψ(CH ₂ NH)I-	G-	I-	L-	T-	V-OH
	Ψ (4-5)	H-A-	A	G-	I	Ψ(CH ₂ NH)G-	I-	L-	T-	V-OH
	Ψ (5-6)	H-A-	A	G-	I-	G	Ψ(CH ₂ NH)I-	L-	T-	V-OH
	Ψ (6-7)	H-A-	A	G-	I-	G	I	Ψ(CH ₂ NH)L-	T-	V-OH
	Ψ (7-8)	H-A-	A	G-	I-	G	I-	L	Ψ(CH ₂ NH)T-	V-OH
15	Ψ (8-9)	H-A-	A	G-	I-	G	I-	L-	T	Ψ(CH ₂ NH)V-OH

These peptide analogues ψ (1-2), ψ (2-3), ψ (3-4), ψ (4-5), ψ (5-6), ψ (6-7), ψ (7-8) and ψ (8-9), also referred to as reduced peptide analogues of Mart-1 27-35, correspond to the latter peptide in which the -CO-NH- bond located between the residues P1 and P2, P2 and P3, P3 and P4, P4 and P5, P5 and P6, P6 and P7, P7 and P8, and P8 and P9 is replaced, in each case, with a -CH₂-NH- bond.

The invention also relates to the peptide analogues of the parent peptide Mart-1 27-35 of melanoma, comprising a bond of the β -homologation type and corresponding to the following formulae :





Sequences


		P1	P2	P3	P4	P5	P6	P7	P8	P9
30	MARTI 27-35	H-A	A-	G-	I-	G-	I-	L-	T-	V-OH
	β1	H-β-homoA	A-	G-	I-	G-	I-	L-	T-	V-OH
	β2	H-A	-β-homoA	G-	I-	G-	I-	L-	T-	V-OH
	β3	H-A-	A	-β-homoG-	I-	G-	I-	L-	T-	V-OH
	β4	H-A-	A-	G	-β-homoI-	G-	I-	L-	T-	V-OH
35	β5	H-A-	A-	G-	I	-β-homoG-	I-	L-	T-	V-OH
	β6	H-A-	A-	G-	I-	G	-β-homoI-	L-	T-	V-OH
	β7	H-A-	A-	G-	I-	G-	I	-β-homoL-	T-	V-OH
	β8	H-A-	A-	G-	I-	G-	I-	L	-β-homoT-	V-OH
	β9	H-A-	A-	G-	I-	G-	I-	L-	T	-β-homoV-OH

These β 1 to β 9 peptide analogues of Mart-1 27-35 correspond to the latter peptide in which the -CO-NH- bond located between the residues P1 and P2, P2 and P3, P3 and P4, P4 and P5, P5 and P6, P6 and P7, P7 and P8, and P8 and P9, is replaced, in each case, with an -CH₂-CO-NH- bond.

The invention also relates to the peptide analogues of the mutated parent peptide Mart-1 27-35 Leu²⁸, corresponding to the peptide Mart-1 27-35 in which the alanine in position 28 is replaced with leucine, and comprising a bond of the ###-homologation type between the same residues as in the case described above for the peptide analogues β 1 to β 9.

The invention is also the peptide analogues of MART1 27-35 in which at least one of the -CO-NH- bonds is replaced with a -CH₂-CH₂- bond, such as the analogues 1 to 8 below (also referred to as carba pseudopeptide analogues of MART):





Séquences

	P1	P2	P3	P4	P5	P6	P7	P8	P9
M58-66	H-G	-L	-L	-G	-F	-V	-F	-T	-L-OH
Ψ(1-2)	H-G	Ψ(CH ₂ NH)L	-L	-G	-F	-V	-F	-T	-L-OH
Ψ(2-3)	H-G	-L	Ψ(CH ₂ NH)L	-G	-F	-V	-F	-T	-L-OH
Ψ(3-4)	H-G	-L	-L	Ψ(CH ₂ NH)G	-F	-V	-F	-T	-L-OH
Ψ(4-5)	H-G	-L	-L	-G	Ψ(CH ₂ NH)F	-V	-F	-T	-L-OH
Ψ(5-6)	H-G	-L	-L	-G	-F	Ψ(CH ₂ NH)V	-F	-T	-L-OH
Ψ(6-7)	H-G	-L	-L	-G	-F	-V	Ψ(CH ₂ NH)F	-T	-L-OH
Ψ(7-8)	H-G	-L	-L	-G	-F	-V	-F	Ψ(CH ₂ NH)T	-L-OH
Ψ(8-9)	H-G	-L	-L	-G	-F	-V	-F	-T	Ψ(CH ₂ NH)L-OH

These peptide analogues ψ (1-2), ψ (2-3), ψ (3-4), ψ (4-5), ψ (5-6), ψ (6-7), ψ (7-8) and ψ (8-9), also referred to as reduced analogues of M58-66, correspond to the latter peptide in which the -CO-NH- bond located between the residues P1 and P2, P2 and P3, P3 and P4, P4 and P5, P5 and P6, P6 and P7, P7 and P8, and P8 and P9 is replaced, in each case, with a -CH₂-NH- bond.

The invention also relates to the peptide analogues of parent peptides of the AIDS virus, in particular of the peptides NEF 84-92 and GAG 77-85, the said peptide analogues corresponding to the said parent peptides in which at least one of the -CO-NH- peptide bonds is modified, with the exception of modifications of the retro or retro-inverso type, the said analogues being chosen in particular from :

- those in which at least one of the -CO-NH- bonds is replaced with a -CH₂-NH-, such as the analogues NEFRD1-8 of the peptide NEF, and GAGRD1-8 of the peptide GAG, below:

			Purity
15	NEF	AVDLSHFLK	98%
	NEFRD1	A Ψ (CH ₂ -NH)VDLSHFLK	99%
	NEFRD2	AV Ψ (CH ₂ -NH)DLSSHFLK	93%
	NEFRD3	AVD Ψ (CH ₂ -NH)LSHFLK	97%
	NEFRD4	AVDL Ψ (CH ₂ -NH)SHFLK	98%
20	NEFRD5	AVDLS Ψ (CH ₂ -NH)HFLK	86%
	NEFRD6	AVDLSH Ψ (CH ₂ -NH)FLK	98%
	NEFRD7	AVDLSHF Ψ (CH ₂ -NH)LK	92%
	NEFRD8	AVDLSHFL Ψ (CH ₂ -NH)K	91%
25			Purity
	GAG	SLYNTVATL	96%
	GAGRD1	S Ψ (CH ₂ -NH)LYNTVATL	95%
	GAGRD2	SL Ψ (CH ₂ -NH)YNTVATL	95%
	GAGRD3	SLY Ψ (CH ₂ -NH)NTVATL	95%
30	GAGRD4	SLYN Ψ (CH ₂ -NH)TVATL	95%
	GAGRD5	SLYNT Ψ (CH ₂ -NH)VATL	95%
	GAGRD6	SLYNTV Ψ (CH ₂ -NH)ATL	95%
	GAGRD7	SLYNTVA Ψ (CH ₂ -NH)TL	95%
	GAGRD8	SLYNTVAT Ψ (CH ₂ -NH)L	95%


- those in which at least one of the -CO-NH- bonds is replaced with a -CHOH-NH- bond, such as the following analogues NEFHEA1-8 of the peptide NEF:

	NEF	AVDLSHFLK	Purity	Mass
	NEFHEA1	AΨ(CHOH-NH)VDLSHFLK	98%	1046.50
	NEFHEA2	AVΨ(CHOH-NH)DLSHFLK	95%	1046.56
5	NEFHEA3	AVDΨ(CHOH-NH)LSHFLK	83%+15%	1029.00
	NEFHEA4	AVDLΨ(CHOH-NH)SHFLK	98%	1047.97
	NEFHEA5	AVDLSΨ(CHOH-NH)HFLK	99%	1047.08
	NEFHEA6	AVDLSHΨ(CHOH-NH)FLK		
	NEFHEA7	AVDLSHFΨ(CHOH-NH)LK	97%	1047.19
10	NEFHEA8	AVDLSHFLΨ(CHOH-NH)K	97%	1046.77

Peptide analogues which are preferred in the context of the present invention are characterized in that at least one of the amino acids in the peptide chain of the parent peptide is substituted with a non-protein-generating amino acid as defined above.




Advantageously, the non-protein-generating amino acids used are chosen from the following amino acids:

- the amino acids of D configuration,
- the α-amino acids of the general formula :




in which :

- R₁, R₂ and R₃, represent, independently of each other : a hydrogen atom, a hydroxyl, an alkyl radical of 1 to 25 carbon atoms, a radical containing an allyl group and having from 3 to 25 carbon atoms, a radical containing one or more aromatic or non-aromatic rings, in particular an aryl group, and having from 6 to 25 carbon atoms, and in particular the following groups: -CH₃ (methyl), -CH₂CH₃ (ethyl), -(CH₂)₄-CH₃, -CH(CH₃)₂ (isopropyl), -C(CH₃)₃ (tert-butyl), -Φ (phenyl), -CH₂ Φ (benzyl), -CH₂ Φ Cl (para-chlorobenzyl), -CH₂-CH₂ Φ (2-phenylethyl), -CH₂CHCH₂ (alkyl), methylfluorenyl, -CH₂CONH₂ (glycolamide), -CH₂CON Φ₂ (benzhydrylglycolamide), -CHOHΦ,




it being understood that one of the two groups R_2 and R_3 can represent a side chain of natural amino acids when either R_1 or the other of the two groups R_2 and R_3 do not represent a hydrogen atom,

- where appropriate, R_1 , R_2 , C_α and N form an aromatic or non-aromatic heterocycle of 4 to 8 carbon atoms, which may be substituted, in particular a heterocycle of the formula :




- the β -amino acids of the general formula :



in which R_1 , R_2 and R_3 , independently of each other, represent a side chain of a natural amino acid or are as defined above,


in particular :

. the β -homo amino acids of the formula:




in which R_1 and R_2 are as defined above, or

. the α -hydroxy β -homo amino acids of the formula :




in which R_1 and R_2 are as defined above,

- the γ -amino acids of the general formula:



in which R_1 , R_2 , R_3 and R_4 represent, independently of each other, a side chain of a natural amino acid, or R_1 , R_2 and R_3 , are as defined above and R_4 has the same meaning as that given above for R_1 , R_2 and R_3 ,

in particular the statin derivatives of the formula :






5




in which R_1 and R_2 are as defined above.

Advantageously, the non-protein-generating amino acids used in the context of the present invention are chosen from;




10






15



20



25




30


More particularly the invention relates to the peptide analogues of the parent peptides of influenza virus, in particular the parent peptide M58-66, the said peptide analogues corresponding to the said parent peptides in which at least one of the amino acids of the peptide chain is substituted with a non-protein-generating amino acid, such as the following analogues:

35


[Tic62]M58-66




[3-Pya62]M58-66




[2-Tha62]M58-66




[D-Phe62]M58-66



[Tic64]M58-66




[3-Pya64]M58-66




16


[2-Tha64]M58-66




[D-Phe64]M58-66




[Cha62]M58-66




[N-MePhe62]M58-66






[pCl-Phe62]M58-66



[Phg62]M58-66



17



Advantageously, the abovementioned peptide analogues of the invention are selected from those which are capable:

- on the one hand, of being recognized by the molecules of the MHC and of associating with these molecule, in particular by carrying out the following method :

incubation (in particular for about 2 hours at 25°C and then for about 12 hours at 4°C) of the peptide analogue in the presence of molecules of the MHC, derived from the lysis of human or animal cells, or purified in particular by affinity chromatography from human or animal cell lines, on a solid support coated with a first antibody, in particular a monoclonal antibody, which specifically recognizes the molecules of the MHC in their conformation which is dependent on their binding to the said peptide analogue

addition to the above solid support of a second antibody which is labelled, in particular by means of coupling with a radioactive, enzymatic or fluorescent label, the said labelled antibody specifically recognizing either the molecules of the MHC in their conformation which is dependent on their binding to the peptide analogue, or a molecule which itself binds specifically to the molecules of the MHC in their abovementioned conformation, in particular β 2-microglobulin which specifically recognizes the molecules of the MHC of category I,

detection, after rinsing the solid support, of the possible presence of the second labelled antibody which has remained bound to the solid support, thereby demonstrating an effect of recognition and association between the molecules of the MHC and the peptide analogue studied,

5 - and, on the other hand, of forming a complex with the said molecules of the MHC, the stability of which complex can be evaluated by carrying out a method for monitoring over time the binding established between the peptide analogue and the molecules of the MHC, this method advantageously being carried out according to a protocol which is identical to the above method, but in which the step of
10 incubation of the peptide analogue in the presence of the molecules of the MHC on the solid support coated with the said first antibody is carried out (advantageously at a temperature of 37°C) for times ranging from a few minutes to several days.

The peptide analogues of the invention must be recognized by the molecules of the MHC and must associate with these molecules, in particular in the context of
15 carrying out the recognition test described above. This association can be weak (detectable at peptide analogue concentrations of the order of 10^{-4} to 10^{-5} M), intermediate (detectable at peptide analogue concentrations of the order of 10^{-5} to 10^{-7} M), or strong (detectable at peptide analogue concentrations of the order of 10^{-8} to 10^{-9} M).

20 The peptide analogues recognized by the molecules of MHC in the context of the present invention are preferably capable of binding for at least about 30 minutes to the said molecules of the MHC.

More particularly, the invention relates to the peptide analogues as described above and characterized in that they are selected from those which are capable :

25 - of inducing *in vitro* the appearance and growth of cytotoxic T lymphocytes from human or animal cells, in particular from peripheral blood mononuclear cells (PBMCs) in the presence of factors required for the growth and differentiation of the cytotoxic T cells,

30 - of inducing *in vitro* cytolysis, by means of cytotoxic T lymphocytes, of target cells carrying at their surface the peptide analogue associated with the molecules of the MHC, the said cytotoxic T lymphocytes advantageously being taken from a patient suffering from a pathology in which the parent peptide of the peptide analogue studied is involved,

35 - and of inducing *in vitro* the secretion of cytokines (or interleukins) by means of the abovementioned cytotoxic T lymphocytes, in particular IL-2, IL-4 or γ -interferon,

the said peptide analogues thus selected being :

. either receptor agonists (TCR) which recognize the antigen (i.e. the parent peptide) of the cytotoxic T cells, and are derived from parent peptides which themselves behave as agonists or antagonists of the said receptors,

. or partial agonists of the said receptors, and are derived from parent peptides which themselves behave as agonists of the said receptors, these partial agonists inducing, in particular, the secretion of one or more cytokines other than those whose secretion is induced by means of the parent peptides.

The invention relates more particularly to the peptide analogues as described above and characterized in that they are selected from those :

- which are capable of inducing *in vitro* the appearance and growth of cytotoxic T lymphocytes from human or animal cells, in particular from peripheral blood mononuclear cells (PBMCs), in the presence of factors required for the growth and differentiation of cytotoxic T cells,

- which do not induce *in vitro* the cytolysis, by means of cytotoxic T lymphocytes, of target cells carrying at their surface the peptide analogue associated with the molecules of the MHC, the said cytotoxic T lymphocytes advantageously being taken from a patient suffering from a pathology in which the parent peptide of the peptide analogue studied is involved,

- which do not induce *in vitro* the secretion of cytokines (or interleukins) by means of the abovementioned cytotoxic T lymphocytes, in particular IL-2, IL-4 or γ -interferon,

the said peptide analogues thus selected being antagonists of the cytotoxic T cell receptors.

The invention also relates to the use of peptide analogues as defined above for the preparation of medicinal products, in particular vaccines, intended for preventing or treating pathologies in which the parent peptides are agonists or antagonists of receptors which recognize the antigen of the cytotoxic T cells, and more particularly neurodegenerative pathologies which are infectious (of viral or bacterial origin), tumoural, autoimmune and allergic.

Among the tumoural pathologies which can be treated in the context of the present invention, mention may be made of melanoma, in particular with the aid of the peptide analogues of the peptide MART1 27-35 which are described above.

Among the diseases of viral origin which can be prevented or treated in the context of the present invention, mention may be made of :

- AIDS brought about by the human immunodeficiency virus HIV1 or HIV2, in particular with the aid of peptide analogues of the peptides NEF and GAG which are described above,

- paraplegia associated with HTVL-1, or T-cell leukemia in adults, brought about by the human T-cell leukemia virus (HTLV virus),

- infections brought about by the syncitial respiratory virus,
- infections brought about by coxsakie virus, for example acute lymphocytic meningitis,
- infections brought about by the Epstein-Barr virus, for example infectious mononucleosis,
- infections brought about by cytomegalovirus, for example cytomegalic inclusion disease,
- herpes brought about by the human herpes virus,
- herpes brought about by virus 6 of herpes simplex,
- infections brought about by human B19 parvovirus, for example infectious gastro-enteritis,
- hepatitis B brought about by the hepatitis B virus,
- hepatitis C brought about by the hepatitis C virus,
- influenza brought about by the influenza virus, in particular with the aid of peptide analogues of the peptide M58-66 which are described above,
- German measles brought about by the rubella virus,
- infections brought about by the Dengue virus, for example arboviruses,
- colds, rhinitis, and coryza brought about by rhinoviruses,
- aphtous fever brought about by the aphtous fever virus,
- cervical cancer brought about by the pillomavirus (HPV).

Among the main autoimmune diseases which can be treated in the context of the present invention, mention may be made of those collated in Table A below.

Table A

Main autoimmune diseases (from top to bottom, autoimmune diseases of specific organs to autoimmune diseases not of specific organs).

5	Disease	Autoantigen involved
	Hashimoto's thyroiditis	Thyroglobulin, microsomes
	Basedow's disease	TSH receptor
	Addison's disease	Corticoadrenal
10	Hypophyseal insufficiency	Pituitary
	Biermer's gastritis	Parietal cell of the stomach
		Intrinsic factor
	Certain sterilities	Spermatozoa, ovaries
15	Type 1 juvenile diabetes	Islets of Langerhans, insulin
	Goodpasture's syndrome	Glomerular basal membrane
	Myasthenia	Striated muscle, acetylcholine receptor
20	Acute articular rheumatism	Myocardium (streptococci)
	Pemphigus	Epidermal intercellular bridges
	Bullous pemphigoid	Cutaneous basal membrane
	Herpetiform dermatitis	Gliadin, reticulin
	Vitiligo	Melanocytes
25	Pelade	Hair follicle
	Psoriasis	
	Sympathic ophthalmia	Uvea
	Uveitis	Anterior chamber of the eye
	Guillain-Barré's syndrome	
30	Multiple sclerosis	Myelin
	Haemolytic anaemia	Red blood cells
	Idiopathic thrombopaenic	
	Purpura	Platelets
	Idiopathic leukopaenia	Granulocytes
35	Primitive biliary cirrhosis	Mitochondria
	Active chronic hepatitis	Smooth muscle, nuclei
	Haemorrhagic rectocolitis	
	Crohn's ileitis	Colon (<i>E. coli</i>)
	Gougerot-Sjögren's syndrome	Nuclei: SS-A, SS-B

	Rheumatoid arthritis	IgG, nuclei
	Dermatopolymyositis	Nuclei: Jo1, muscles
	Scleroderma	Nuclei: Scl-70
	Mixed connectivty	Nuclei: RNP
5	Discoid lupus erythematosus	Nuclei:
	Disseminated lupus erythematosus	Nuclei: DNA, Sm antigen
		Clotting factors
		Cardiolipin, etc.

10

The invention also relates to any pharmaceutical composition, characterized in that it comprises, as active principle, at least one peptide analogue (agonist, which may be a partial agonist, or antagonist) as defined above, in combination with a pharmaceutically acceptable vehicle.

15

Advantageously, the pharmaceutical compositions according to the invention are in a form which can be administered orally or parenterally, in particular at a rate of about 500 µg to 5 mg at a time, in particular at a rate of 3 times a day.

20

More particularly, the invention relates to the pharmaceutical compositions as described above, containing, as active principle, at least one antagonist according to the invention, and their use in the context of treating autoimmune diseases.

25

The invention also relates to any vaccine, characterized in that it comprises, as active principle, at least one peptide analogue, preferably an agonist peptide analogue, which may be a partial agonist, as defined above, in combination with a pharmaceutically acceptable vehicle.

30

Advantageously, the vaccines according to the invention are in a form which can be administered orally or parenterally, in particular at a rate of about 500 µg to 5 mg at a time, in particular at a rate of 3 times a day.

35

The invention also relates to any composition intended for *in vivo* diagnosis of abovementioned pathologies involving the immune response to cell mediation, in particular the cytotoxic T lymphocytes, or to the *in vivo* evaluation of the immune response in the context of the said pathologies, by carrying out a skin hypersensitivity reaction by means of intradermal injection of the said diagnostic composition, characterized in that it comprises at least one peptide analogue, preferably an agonist peptide analogue, which may be a partial agonist, as defined above, in combination with a biologically acceptable vehicle.

The invention also relates to the use of peptide analogues as described above for the preparation of the said compositions intended for the *in vivo* diagnosis of the abovementioned pathologies.

The invention also relates to complexes between a peptide analogue as defined
5 above and a molecule of the major histocompatibility complex (also referred to as MHC-peptide analogue binary complexes).

The immune response involves the recognition of an endogenous or exogenous antigen by specialized cells. In order to be recognized, the antigen must first be presented in a suitable manner by antigen presenting cells (APCs). Whereas the B
10 lymphocytes recognize epitopes borne by the unmodified intact antigens, the presentation of the antigen to the T lymphocytes is more complex since the antigen is first internalized by the presenting cell, proteolysed and then possibly re-expressed at its surface in the form of peptide fragments in association with the proteins of the major histocompatibility complex (MHC). The T lymphocyte, which does not
15 recognize the native antigen, recognizes a peptide fragment associated with an MHC molecule.

These MHC molecules belong essentially to two categories: I and II.

The molecules of category I are transmembrane glycoproteins consisting of a heavy polymorphic α chain associated non-covalently with a light non-glycosylated
20 β_2m chain. Their crystallographic structure has been resolved (Bjorkman et al. (1987), Nature, 329: 506-512), and shows the presence of a channel forming the presentation site of the peptide, the base of which is composed of eight β sheets and the edges are composed of two α helices. These molecules are presented at the surface of virtually all cells.

The molecules of category II are also membrane glycoproteins consisting of two polymorphic chains α et β which are non-covalently linked to form, as shown in the crystallographic structure recently elucidated (Brown et al. (1993), Nature, 364:
25 33-39), a β -pleated platform supporting two α helices. The channel formed is the presentation site of the peptide. These molecules are expressed only at the surface of certain cells, including macrophages and B cells.

The cytotoxic T lymphocytes (cells possessing CD8 labels) recognize proteolytic fragments of viral proteins associated with the MHC molecules of category I and bring about the lysis of cells bearing the antigen.

The helper T lymphocytes (cells bearing CD4 labels) recognize fragments of
35 exogenous proteins captured by endocytosis which are presented in association with the MHC molecules of category II and induce the cellular stimulation of the immune response.

The invention also relates to complexes between a peptide analogue according to the invention and a T cell receptor (which are also referred to as T receptor-peptide analogue complexes).

5 The invention is also directed towards complexes between a molecule of the major histocompatibility complex, a peptide analogue as defined above and a T cell receptor (which is also referred to as an MHC-peptide analogue-T receptor ternary complex).

The invention also relates to the use of peptide analogues as defined above, for carrying out a method for the *in vitro* diagnosis of the pathologies mentioned above.

10 More particularly, the invention relates to any method for the *in vitro* diagnosis of pathologies involving the immune response to cell mediation, in particular the cytotoxic T lymphocytes, i.e. pathologies associated with the presence, in a patient's body, of exogenous or endogenous peptides which can interact with molecules of the MHC, and which are liable to be directly or indirectly involved in the process of
15 development of these pathologies in man or animals, characterized in that it comprises:

- placing a biological sample obtained from a patient, in particular the blood or any biological sample liable to contain lymphocytes, in contact with a peptide analogue as defined above, under conditions which allow reaction between the
20 receptors of the T cells liable to be present in the biological sample, and the abovementioned binary complex formed between the said peptide analogue and the molecules of the MHC present in the said sample;

- the *in vitro* detection of the MHC-peptide analogue-T receptor ternary complex which may be formed in the preceding step.

25 The abovementioned diagnostic methods of the invention are advantageously carried out in the following way:

- incubation of the said biological sample with peptide analogues according to the invention, the said peptide analogues being bound to a solid support, in particular to the inside of the wells of microtitration plates of the type usually used for carrying
30 out detection or assay techniques which are well known under the name ELISA (Enzyme Linked Immuno Sorbent Assay),

- rinsing of the solid support,

- incubation of the components which have remained bound to the solid support after the preceding rinsing step, with a medium containing antibodies, in
35 particular antibodies raised against the ternary complex according to the invention, which are labelled (in particular by radioactive, enzymatic or fluorescent means) or which are capable of being recognized in turn by a labelled reagent,

- rinsing of the solid support,

- detection of the labelled antibodies which have respectively remained bound to the ternary complexes during the preceding incubation step.

A subject of the invention is also equipment or kits for carrying out the methods of *in vitro* diagnosis as described above, comprising:

5 - a peptide analogue as defined above;
 - reagents for making a medium suitable for forming an immunological reaction;

10 - reagents for detecting the ternary complex according to the invention, which has been produced after the immunological reaction, the said reagents optionally containing a label or being capable of being recognized in turn by a labelled reagent, more particularly in the case in which the peptide analogue is not labelled.

15 The invention also relates to antibodies directed against the MHC-peptide analogue binary complexes as defined above, the said antibodies being as obtained by immunizing an animal with at least one of the abovementioned complexes, the said antibodies being capable of forming a complex with these binary complexes.

 The antibodies according to the invention are monoclonal or polyclonal antibodies.

20 The abovementioned polyclonal antibodies are obtained by immunizing an animal with at least one MHC-peptide analogue complex according to the invention, followed by recovery of the desired antibodies in purified form, by taking the serum of the said animal and separating out the antibodies from the other constituents of the serum, in particular by affinity chromatography on a column onto which is bound an antigen which is specifically recognized by the antibodies, in particular an MHC-peptide analogue complex according to the invention.

25 The monoclonal antibodies according to the invention can be obtained by the hybridoma technique, the general principle of which is recalled below.

30 In a first stage, an animal, generally a mouse (or cells under culture in the context of *in vitro* immunizations), is immunized with an MHC-peptide analogue complex according to the invention, the B lymphocytes of which are then capable of producing antibodies against the said complex. These antibody-producing lymphocytes are then fused with (in particular mouse) "immortal" myeloma cells to give rise to hybridomas. Using the heterogeneous mixture of cells thus obtained, a selection is then made of cells capable of producing a particular antibody and of multiplying indefinitely. Each hybridoma is multiplied in the form of a clone, each
35 leading to the production of a monoclonal antibody whose properties of recognition with respect to the MHC-peptide analogue complex of the invention may be tested by ELISA, for example, by one- or two-dimensional immunotransfer, by immunofluorescence, or with the aid of a biosensor. The monoclonal antibodies thus

selected are then purified, in particular according to the affinity chromatography technique described above.

The antibodies according to the invention are more particularly characterized in that they are capable of forming a complex with MHC-peptide analogue complexes and/or with complexes of MHC-parent peptide or protein correspondant to the said peptide analogues.

Advantageously, the antibodies raised against the MHC-peptide analogue complexes of the invention recognize the abovementioned complexes of MHC-parent peptide or protein, with an affinity which is at least equal to that presented by the anti(MHC-parent peptide or protein complex) antibodies with respect to the MHC-parent peptide or protein complexes.

The affinity which is concerned above can be measured by means of the affinity constant at equilibrium, K_a , of the complexes involving one of the abovementioned antibodies with one of the abovementioned complexes.

The invention also relates to any pharmaceutical composition, characterized in that it comprises antibodies as defined above, in combination with a physiologically acceptable vehicle, as well as their use in the context of treating the abovementioned pathologies.

The invention also relates to the process for screening peptide analogues as defined above, characterized in that it comprises the following steps :

- . incubation (in particular at a temperature of 37°C), for times ranging from a few minutes to several days, of the peptide analogue in the presence of molecules of the MHC, derived from the lysis of human or animal cells, or purified in particular by affinity chromatography from human or animal cell lines, on a solid support coated with a first antibody, in particular a monoclonal antibody, which specifically recognizes the molecules of the MHC in their conformation which is dependent on their binding to the said peptide,

- . addition to the preceding solid support of a second antibody which is labelled, in particular by means of coupling with a radioactive, enzymatic or fluorescent label, the said labelled antibody specifically recognizing either the molecules of the MHC in their conformation which is dependent on their binding to the peptide analogue, or a molecule which itself binds specifically to the molecules of the MHC in their abovementioned conformation, in particular β 2-microglobulin which specifically recognizes the molecules of the MHC of category I,

- . detection, after rinsing the solid support, of the possible presence of the second labelled antibody which has remained bound to the solid support,

- . evaluation of the duration of the association between the said peptide analogue and the molecules of the MHC.

The invention also relates to any set or kit for carrying out a process for screening peptide analogues as defined above, comprising:

- molecules of the MHC, and/or
- antibodies which specifically recognize the molecules of the MHC in their conformation which is dependent on their binding to the said peptide analogue, which antibodies are advantageously bound to a solid support, or are supplied with the reagents required for binding them to the solid support, and/or
- antibodies which are labelled, in particular by means of coupling with a radioactive, enzymatic or fluorescent label, these antibodies specifically recognizing either the molecules of the MHC in their conformation which is dependent on their binding to the peptide analogue, or a molecule which itself binds specifically to the molecules of the MHC in their abovementioned conformation, in particular β 2-microglobulin which specifically recognizes the molecules of the MHC of category I, and/or
- a protocol for carrying out the said process, and/or
- a control peptide.

The invention will be further illustrated with the aid of the following examples of the production and study of the peptide analogues as described above.

I- Synthesis of the peptide analogues of the invention :

A) General features :

By way of illustration, the liquid-phase synthesis method consists in successively condensing the aminoacyls two by two in the required order, or in condensing aminoacyls and fragments which have been formed beforehand and which already contain several aminoacyls in the appropriate order, or alternatively several fragments thus prepared beforehand, it being understood that care will have been taken to protect beforehand all the reactive functions borne by these aminoacyls or fragments, with the exception of the amine functions of one and carboxyl of the other or vice-versa, which should normally be involved in the formation of the peptide bonds, in particular after activation of the carboxyl function, according to the methods which are well known in peptide synthesis.

It will be possible, for example, to use protecting groups of urethane type (Boc, Fmoc benzyloxycarbonyl or allyloxycarbonyl) to protect the N-terminal ends of the amino acids, and groups of ester type (methyl, ethyl, benzyl, tert-butyl, allyl or alternatively benzhydrylglycolamide) to protect the C-terminal ends of the amino acids.

Such a synthesis can be carried out by first condensing the aminoacyl residue AA1, whose COOH function is protected, with the aminoacyl residue AA2 whose NH₂ function is protected. The amine function of the residue AA2 in the fragment AA2-AA1 thus obtained is then deprotected, in order subsequently to condense the said fragment with the aminoacyl residue AA3 whose amine function is protected. The preceding steps are repeated as many times as there are aminoacyl residues to be introduced into the chain of the retro analogues to be synthesized.

According to another preferred method of the invention, use is made of the technique described by R.D. Merrifield in the article entitled "Solid phase peptide synthesis" (J. Am. Chem. Soc. (1963), 85, 2149-2154).

In order to manufacture a peptide chain according to the Merrifield process, use is made of a highly porous polymer resin, onto which is bound the first C-terminal amino acid (in this instance AA1-OH) of the chain. This amino acid is bound to the resin via its carboxyl group and its amine function is protected, for example with the t-butoxycarbonyl group.

When the first C-terminal amino acid is thus bound to the resin, the protecting group is removed from the amine function by washing the resin with an acid.

The amino acids which will make up the peptide chain are thus bound, one after the other, to the amino group each time the portion of the peptide chain which is already formed, and which is attached to the resin, undergoes preliminary deprotection.

When all of the desired peptide chain is formed, the protecting groups are removed from the various amino acids constituting the peptide chain and the peptide is detached from the resin, for example using hydrofluoric acid.

B) Particular case of the peptide analogues of the invention :

One or more of the synthesis steps described above can be interrupted in order to insert a bond other than the -CO-NH- bond between certain aminoacyl residues, and /or one or more non-protein-generating amino acids.


B.1.) Synthesis of peptide analogues containing modified peptide bonds :

The notation ψ (Spatola & Darlak, 1988) is generally used to denote the pseudopeptide bond which is replacing the amide bond (CONH). Among the modifications most frequently encountered, mention may be made, for example, of the following, the syntheses of which are detailed in the bibliographic references given : the methyleneamino (reduced) bond $\psi[\text{CH}_2\text{NH}]$ (Szelke *et al.*, 1982 ; Martinez *et al.*, 1985 ; Sasaki & Coy, 1987) ; retro-inverso bond $\psi[\text{NHCO}]$ (Shemyakin *et al.*, 1969 ; Goodman & Chorev, 1979 ; Chorev & Goodman, 1993);

olefin bond $\psi[\text{CH}=\text{CH}]$ (Hann *et al.*, 1982 ; Kempf *et al.*, 1991 ; Bohnstedt *et al.*, 1993) ; carba bond $\psi[\text{CH}_2\text{CH}_2]$ (Rodriguez *et al.*, 1990a, 1990b) ; methyleneoxy (ether) bond $\psi[\text{CH}_2\text{O}]$ (TenBrink, 1986 ; Breton *et al.*, 1990) ; ketomethylene bond $\psi[\text{COCH}_2]$ (Harbeson & Rich, 1989 ; Gonzalez-Muniz *et al.*, 1995), hydroxyethylene bond $\psi[\text{CH}(\text{OH})\text{CH}_2]$ (Evans *et al.*, 1985 ; Wuts *et al.*, 1992).

1) Synthesis of the reduced pseudopeptides corresponding to the sequences of MART and of the matrix peptide M58-66 :


The contribution of the amine function of each peptide bond of the antigen Mart-1 27-35 (AAGIGILTV) of the protein Mart-1 du melanoma and of the antigen M58-66 (GLLG FVFTL) of the matrix protein of influenza virus was evaluated as regards the interactions of these antigens with their receptors: the molecule of category I of the MHC HLA-A2 and TCR.



Scheme 1: Schematic representation of the peptide bonds and reduced bonds.


The reduced bond, or methyleneamino bond, has the particular feature of being more flexible than the peptide bond on account of a free rotation of the carbon-carbon bond (scheme 1). The introduction of a reduced bond can locally modify the orientation of the amino acid side chains. The reduced bond can exist in its protonated form at physiological pH. The changes brought about by the presence of a reduced bond, i.e. the change in the structure and the change in the hydrophilic nature of the parent peptide, can have implications in the phenomena of binding and recognition of the antigen.

The synthesis of each reduced peptide chain was performed in solid phase and by standard Fmoc chemistry. The reduced bond is obtained by condensing an N-protected α -amino aldehyde with the α -amino acid immobilized on the resin. The imine form is reduced *in situ* with sodium cyanoborohydride to give the reduced bond according to the method described by Sasaki and Coy in 1987 (scheme 2).



10 Scheme 2: Incorporation of the reduced bond in solid phase. PG: Fmoc protecting group.

The α -amino aldehyde is synthesized by the method of Fehrentz and Castro, 1983. This method allows the rapid production, without racemization, of the N-protected α -amino aldehyde from the N,O-dimethylhydroxamate of the corresponding N-protected amino acid (scheme 3).



25 Scheme 3: Synthesis of N-protected α -amino aldehydes by the method of Fehrentz and Castro. PG: Fmoc protecting group.

30 The following amino acids of the sequence of the reduced peptide are coupled in a conventional manner. After synthesis, the reduced peptide is deprotected and detached from the resin with trifluoroacetic acid (TFA). Each reduced peptide is purified by reverse-phase high pressure liquid chromatography (RP-HPLC). The mass of each peptide is controlled by matrix assisted laser desorption (MALDI) with the aid of a Bruker Protein TOF spectrometer.

35 The retention times (Tr) on an RP-HPLC C18 column in a gradient of from 5 to 65 % B over 30 min (with A = water containing 0.1 % TFA and B = acetonitrile (ACN) containing 0.08 % TFA) and the masses of the various peptides are given in Tables 1 et 2.

Table 1: Sequences and characteristics of the reduced peptide analogues of Mart-1 27-35

		Sequences									Tr	mass	
		P1	P2	P3	P4	P5	P6	P7	P8	P9			
5													
	MART1 27-35	H- A	- A	- G	- I	- G	- I	- L	- T	- V	-OH	12.74	814.4
	Ψ(1-2)	H-AΨ(CH ₂ NH)A	-G	- I	- G	- I	- L	- T	- V	-OH		12.53	802.7
	Ψ(2-3)	H- A- AΨ(CH ₂ NH)G	-I	- G	- I	- L	- T	- V	-OH			12.58	802.9
10	Ψ(3-4)	H- A	- A	-GΨ(CH ₂ NH)I	- G	- I	- L	- T	- V	-OH		11.85	801.6
	Ψ(4-5)	H- A	- A	- G	-IΨ(CH ₂ NH)G	- I	- L	- T	- V	-OH		12.03	801.5
	Ψ(5-6)	H- A	- A	- G	- I	-GΨ(CH ₂ NH)I	- L	- T	- V	-OH		11.29	801.4
	Ψ(6-7)	H- A	- A	- G	- I	- G	-IΨ(CH ₂ NH)L	- T	- V	-OH		12.00	800.4
	Ψ(7-8)	H- A	- A	- G	- I	- G	- I	- L	Ψ(CH ₂ NH)T	-V	-OH	12.77	801.5
15	Ψ(8-9)	H- A	- A	- G	- I	- G	- I	- L	- T	Ψ(CH ₂ NH)V	-OH	12.46	801.3


Table 2: Sequences and characteristics of the reduced peptide analogues of M58-66

		Sequences									Tr	mass
		P1	P2	P3	P4	P5	P6	P7	P8	P9		
20												
	M58-66	H- G	- L	- L	- G	- F	- V	- F	- T	- L	-OH	12.35* 968.0
	Ψ(1-2)	H-GΨ(CH ₂ NH)L	- L	- G	- F	- V	- F	- T	- L	-OH		10.67* 956.2
	Ψ(2-3)	H- G- LΨ(CH ₂ NH)L	- G	- F	- V	- F	- T	- L	-OH			11.18* 957.1
25	Ψ(3-4)	H- G	- L	- LΨ(CH ₂ NH)G	- F	- V	- F	- T	- L	-OH		16.01 957.3
	Ψ(4-5)	H- G	- L	- L- GΨ(CH ₂ NH)F	- V	- F	- T	- L	-OH			15.61 954.3
	Ψ(5-6)	H- G	- L	- L- G	- FΨ(CH ₂ NH)V	- F	- T	- L	-OH			10.80* 955.8
	Ψ(6-7)	H- G	- L	- L- G	- F- VΨ(CH ₂ NH)F	- T	- L	-OH				11.85* 953.2
	Ψ(7-8)	H- G	- L	- L- G	- F- V- FΨ(CH ₂ NH)T	- L	-OH					13.05* 954.7
30	Ψ(8-9)	H- G	- L	- L- G	- F- V- F- TΨ(CH ₂ NH)L	-OH						11.21* 957.3

Retention time in the gradient of from 20 to 80% B over 30 min (with A = water containing 0.1% TFA and B = ACN containing 0.08% TFA).

2) Synthesis of the β analogues of the parent MART peptide and of the mutated (Leu²⁸) MART peptide :

The β analogues are obtained by coupling β -homo amino acids in place of the natural amino acids.




Scheme 4: Schematic representation of an amino acid and of a β -homo amino acid

The β -homo amino acid is an amino acid in which a methylene group has been inserted between the α carbon and the carboxyl. The configuration of the $C\alpha$ is unchanged (scheme 4). The incorporation of a β -homo amino acid in place of an amino acid in a peptide sequence has the consequence of lengthening the peptide chain of the resulting peptide analogues. These longer peptide analogues can be made to twist and brace themselves to penetrate into the binding pocket of the molecule of category I of the MHC. In this case, certain side chains can be displaced laterally or can have their orientation modified, which can result in preferential interactions with the molecule of category I of the MHC and/or TCR.

The β -homo amino acids are obtained by chemical synthesis in several steps starting with the corresponding N-protected amino acids, according to the scheme below (scheme 5).

The amino acid protected on its amine function with a Boc group is, in a first stage, activated by means of the mixed anhydride method. The activated amino acid reacts with diazomethane to give the diazomethyl ketone 2. The Wolff rearrangement of this diazomethyl ketone in methanol, in the presence of silver benzoate and triethylamine, gives the β -homo methyl ester of the amino acid protected with the Boc group. A saponification gives the compound 3. Deprotection of the Boc, followed by reprotection of the amine function with the Fmoc are the final steps for obtaining the β -homo amino acid protected with Fmoc.

Compound 4 is introduced by means of the conventional methods for coupling amino acids in the course of solid-phase synthesis in the Fmoc strategy. In practice, 5 equivalents of the mixture (4/BOP/HOBt) are coupled for 20 min in DMF. The Fmoc group is deprotected with 50% piperidine in DMF and the synthesis is continued conventionally. See the preceding paragraph for the detachment and deprotection of the peptide, as well as for the purification and characterization of the peptides.



Scheme 5: Scheme of synthesis of Fmoc-β-homo amino acids

Tables 3 and 4 give the retention times on RP-HPLC of the ### peptides on a C18 column (gradient of from 5 to 65% B over 30 min with A = water containing 0.1% TFA and B = ACN containing 0.08% TFA) as well as their mass analysis by MALDI.

Table 3: Sequences and characteristics of the β peptide analogues of Mart-1


27-35

		Sequences									Rt	mass	
		P1	P2	P3	P4	P5	P6	P7	P8	P9			
30	MART1 27-35	H- A	- A	- G	- I	- G	- I	- L	- T	- V	-OH	12.74	814.4
	β1	H-β-homoA	- A	- G	- I	- G	- I	- L	- T	- V	-OH	12.62	868.9*
	β2	H- A-β-homoA	- G	- I	- G	- I	- L	- T	- V	-OH	12.50	829.9	
	β3	H- A	- A-β-homoG	- I	- G	- I	- L	- T	- V	-OH	12.48	828.9	
35	β4	H- A	- A	- G-β-homoI	- G	- I	- L	- T	- V	-OH	12.50	829.5	
	β5	H- A	- A	- G	- I-β-homoG	- I	- L	- T	- V	-OH	12.50	828.9	
	β6	H- A	- A	- G	- I	- G-β-homoI	- L	- T	- V	-OH	12.64	829.0	
	β7	H- A	- A	- G	- I	- G	- I-β-homoL	- T	- V	-OH	12.10	828.9	
	β8	H- A	- A	- G	- I	- G	- I	- L-β-homoT	- V	-OH	11.92	829.1	
40	β9	H- A	- A	- G	- I	- G	- I	- L	- T-β-homoV	-OH	12.68	829.2	

* corresponds to M+K⁺

The synthon Fmoc-Ile ψ (CH₂CH₂)Gly-OH (4 in Figure 7) is necessary for the synthesis of the analogue [Ile³⁰ ψ (CH₂CH₂)Gly³¹] Mart-1 27-35. This synthon is obtained by chemical synthesis in several steps (scheme 7).

The dimethylhydroxamate of Boc- β -homole-OH (1) (see the preceding paragraph for the synthesis of the β -homo amino acids) is reduced with lithium aluminium hydride at low temperature (-25°C) in tetrahydrofuran in order to give the corresponding aldehyde (2). The Horner-Emmons reaction in the presence of triethyl phosphonoacetate gives the ethylenic dipeptide 3. Hydrogenation of the double bond, saponification of the ester, and deprotection and re-protection of the amine function with the Fmoc group allow the Fmoc-Ile ψ (CH₂CH₂)Gly-OH (4) to be obtained.



Scheme 7: Scheme of synthesis of Fmoc-Ile³⁰ ψ (CH₂CH₂)Gly³¹-OH (4)

This synthon (4) is incorporated in the solid phase in Fmoc chemistry. Coupling of 5 equivalents of the mixture (synthon/BOP/HOBt) is carried out twice in succession (20 min each time) in DMF. The Fmoc group is deprotected with 50% piperidine in DMF. The following amino acids are coupled conventionally.

The peptide is deprotected and detached from the resin in TFA, precipitated in ether and purified by RP-HPLC.

The pseudopeptide [Ile³⁰ ψ (CH₂CH₂)Gly³¹] Mart-1 27-35 is obtained in a purity of 100%. Its retention time in a gradient of from 5 to 65% B (with A = water containing 0.1% TFA and B=ACN containing 0.08% TFA is 13.49 min. Analysis of the mass of the pseudopeptide by MALDI with the aid of a Bruker Protein TOF spectrometer gives the expected mass $M+H^+$ 800.0.

4) Synthesis of the carba pseudopeptide analogues 1 to 8 of MART1 27-35

The synthesis of these peptides is carried out according to the procedure described in Rodriguez et al., Tetrahedron Letters, 1990, 31, 7319-7322.

The Mart-1 27-35 carba pseudopeptide 1 is obtained in a purity of 90%. Its retention time in a gradient of from 5 to 65% B (with the eluent : A = water containing 0.1% TFA and B = acetonitrile (ACN) containing 0.08% TFA) is 12.98/13.10 min. The analysis of the mass of the pseudopeptide by MALDI using a Bruker Protein TOF spectrometer gives the expected mass $M+H^+$ 799.1.

The Mart-1 27-35 carba pseudopeptide 2 is obtained in a purity of 98%. Its retention time in a gradient of from 5 to 65% B (with the eluent : A = water containing 0.1% TFA and B = acetonitrile (ACN) containing 0.08% TFA) is 12.90 min. The analysis of the mass of the pseudopeptide by MALDI using a Bruker Protein TOF spectrometer gives the expected mass $M+H^+$ 799.16.

The Mart-1 27-35 carba pseudopeptide 3 is obtained in a purity of 96% (3a), 95% (3b). Its retention time in a gradient of from 5 to 65% B (with the eluent : A = water containing 0.1% TFA and B = acetonitrile (ACN) containing 0.08% TFA) is 13.07 (3a), 13.54 (3b) min. The analysis of the mass of the pseudopeptide by MALDI using a Bruker Protein TOF spectrometer gives the expected mass $M+H^+$ 799.13.

The Mart-1 27-35 carba pseudopeptide 4 is obtained in a purity of 100%. Its retention time in a gradient of from 5 to 65% B (with the eluent : A = water containing 0.1% TFA and B = acetonitrile (ACN) containing 0.08% TFA) is 13.49 min. The analysis of the mass of the pseudopeptide by MALDI using a Bruker Protein TOF spectrometer gives the expected mass $M+H^+$ 800.03.

The Mart-1 27-35 carba pseudopeptide 5 is obtained in a purity of 99.7% (5a), 96% (5b). Its retention time in a gradient of from 5 to 65% B (with the eluent : A = water containing 0.1% TFA and B = acetonitrile (ACN) containing 0.08% TFA) is 12.72 (5a), 14.16 (5b) min. The analysis of the mass of the pseudopeptide by MALDI using a Bruker Protein TOF spectrometer gives the expected mass $M+H^+$ 799.1.

The Mart-1 27-35 carba pseudopeptide 6 is obtained in a purity of 86% (6a), 90% (6b). Its retention time in a gradient of from 5 to 65% B (with the eluent : A = water containing 0.1% TFA and B = acetonitrile (ACN) containing 0.08% TFA) is 13.25 (6a), 13.72 (6b) min. The analysis of the mass of the pseudopeptide by MALDI using a Bruker Protein TOF spectrometer gives the expected mass $M+H^+$ 799.16.

The Mart-1 27 carba pseudopeptide 7 has a retention time in a gradient of from 5 to 65% B (with the eluent : A = water containing 0.1% TFA and B = acetonitrile (ACN) containing 0.08% TFA) of 10.20 (7a), 13.61 (7b) min.

The Mart-1 27-35 carba pseudopeptide 8 is obtained in a purity of 99%. Its retention time in a gradient of from 5 to 65% B (with the eluent : A = water containing 0.1% TFA and B = acetonitrile (ACN) containing 0.08% TFA) is 15.24/15.29 min. The analysis of the mass of the pseudopeptide by MALDI using a Bruker Protein TOF spectrometer gives the expected mass $M+H^+$ 798.

B.2.) Synthesis of peptide analogues comprising non-protein-generating amino acids :

Synthesis of the peptide analogue [Tic62] M58-66

Tetrahydroisoquinoline-3-carboxylic acid is a constrained analogue of phenylalanine. It was introduced into position 62 in the sequence of the influenza matrix peptide in place of the natural phenylalanine in order to study the importance of the orientation of the phenyl group in the interaction with the MHC molecule and TCR.


The peptide analogue was synthesized by standard Fmoc chemistry. The derivative Fmoc-L-Tic-OH (sold by the company Néosystem) was introduced into the growing chain using a conventional coupling process (BOP/HOBt/Fmoc-L-Tic-OH) in a five-fold excess for 20 min in DMF. The deprotection of the Fmoc group was carried out in several steps with 50% piperidine in DMF (4 treatments of 30 min each, separated by 3 washes of the resin with DMF). Assembly of the following amino acids posed no particular problems. After standard deprotection with TFA and HPLC purification, the peptide was obtained in a purity of 90%. . The mass measured using a Bruker Protein TOF spectrometer was the expected mass $M+H^+$ 978.9.

The synthesis of the peptide analogue [Tic64]M58-66 is carried out in the same way as previously, introducing tetrahydroisoquinoline-3-carboxylic acid in place of the phenylalanine in position 64.

Still according to the same protocol :


. the analogues 3-Pya62 and 3-Pya64 of M58-66 were obtained by introducing the amino acid of the formula

38



in place of the phenylalanine in positions 62 and 64, respectively ;


. the analogues 2-Tha62 and 2-Tha64 of M58-66 were obtained by introducing the amino acid of the formula



in place of the phenylalanine in positions 62 and 64, respectively ;


. the analogues D-Phe62 and D-Phe64 of M58-66 were obtained by introducing D-phenylalanine in place of the phenylalanine in positions 62 and 64, respectively ;

. the analogues Cha62 and Cha64 of M58-66 were obtained by introducing the amino acid of the formula




in place of the phenylalanine in positions 62 and 64, respectively ;

. the analogue N-MePhe62 of M58-66 was obtained by introducing the amino acid of the formula




in place of the phenylalanine in position 62 ;

. the analogues pCl-Phe62 and pCl-Phe64 of M58-66 were obtained by introducing the amino acid of the formula



in place of the phenylalanine in positions 62 and 64, respectively ;

. the analogues Phg62 and Phg64 of M58-66 were obtained by introducing the amino acid of the formula



in place of the phenylalanine in positions 62 and 64, respectively.

II- Study of the biological properties of the peptide analogues synthesized:

A) Methodology :

1) Detection of the association between the peptides and the histocompatibility molecules

1.1. Equipment

Sources of histocompatibility molecules

They are currently of two main types : mutant cells and purified histocompatibility molecules.

- The mutant cell used is the human T2 cell (DeMars *et al.*, 1985; Ljunggren & Kärre, 1985 ; Salter & Cresswell, 1986) which is a variant of the T1 line produced by fusion of the CEM T lymphoma and of the B lymphoma 721,174. This cell, which lacks peptide carriers, contains heavy chains of molecules of category I free of peptides which will accept exogenous peptides.

- Histocompatibility molecules of category I purified by affinity chromatography from human B cell lines transformed with EBV can also be used. In this case, the endogenous peptides must be removed by means of a treatment with 1.5 M urea and 12.5 mM sodium hydroxide (pH 11.7) for 1 h at 4°C, followed by removing them by means of a desalination column (PD10, Pharmacia). The histocompatibility molecules are immediately put back in the presence of the peptides to be tested in a PBS buffer with 0.05 % Tween 20, 2 mM EDTA, 0.1% NP40 and 6 mM CHAPS, in the presence of 2 µg/ml B2m to facilitate the reassociation (Gnjatic *et al.*, 1995).

Peptides

The peptides tested are used at concentrations ranging from 100 μ M to 0.1 nM.

5 1.2. Assembly protocol

Aliquots of 8×10^5 T2 cells in a volume of 64 μ l, distributed in Eppendorf microfuge tubes, are placed in the presence of a lysis buffer containing 10 mM PBS, pH 7.5, 1% NP40, protease inhibitors (1 mM PMSF, 100 μ M iodoacetamide, 2 μ g/ml aprotinin, 10 μ M leupeptin, 10 μ M pepstatin and 10 μ g/ml trypsin inhibitor). The lysis is carried out in the presence of the peptides to be tested for 30 min or 1 h at 37°C. After elimination of the undissolved material by centrifugation at 15 000 rpm at 4°C, 140 μ l of PBS containing 0.05% of Tween 20, 3 mM of sodium azide, 1 mM PMSF and 10 mg/ml of bovine albumin are added to the supernatant. Each sample is incubated for 20 h at 4°C in 2 wells of a Nunc or Maxisorb type microtitration plate precoated with a monoclonal antibody (Parham & Brodsky, 1981) (10 μ g/ml in PBS) which recognizes the histocompatibility molecules having one (or more) conformation(s) which comply (complies) for the presentation of the peptides and which is (are) similar to that (those) present at the surface of the cells. The plate coated with antibody is presaturated with bovine albumin at 10 mg/ml in PBS-Tween before adding the sample. The second antibody which allows detection of the assembly of the histocompatibility molecules is directed against beta2m. It is coupled either with biotin (NHS-LC biotin, Pierce) or to alkaline phosphatase (P-5521, Sigma) and is incubated at 2 μ g/ml for 1 h at 37°C. In the case of the use of biotin, an incubation for 45 min at 20-25°C with extravidin coupled with alkaline phosphatase (E-2636, Sigma) is carried out. The activity of the alkaline phosphatase is measured using, as substrate, 4-methyl-umbelliferyl phosphate (M-8883, Sigma) at 100 μ M in 50 mM diethanolamine, pH 9.5 with 1 mM $MgCl_2$. The reading is carried out at 340/460 nm using a Millipore 2300 cytofluorimeter.

30

1.3. Controlling the stability of the HLA/peptide complexes

The material used is either purified HLA or the lysate of the T2 cell. With purified HLA, the endogenous proteins need to be removed, as described in paragraph I-1, and the HLA placed in the presence of the peptide to be tested in an Eppendorf tube at 37°C for times ranging from a few minutes to several days. The subsequent phase of incubation on a 96-well plate (see paragraph I-2) with the anti-HLA antibody (Parham & Brodsky, 1981) is carried out for 1 h at 37°C. The detection is standard (Teillaud *et al.*, 1982). With the T2 cell lysate, all the

35

incubations are also carried out at 37°C after all the protease inhibitors have been added.

1.4. Analysis of the conformation of the HLA/peptide complexes by BIAcore surface plasmonic resonance

This machine measures, with the aid of a reflected light beam, the molecular interactions which take place in real time in a continuous flow of buffer. The anti-HLA antibodies are diluted to 25 µg/ml in pH 7.4 sodium acetate buffer and are covalently bound via their free amines to the carboxymethyl dextran matrix of the "Sensor Chip CM5" activated with 50 mM NHS and 200 mM EDC. The injection is carried out at a rate of 5 µl/min for 6 minutes and the residual reactive groups are then blocked with 30 µl of pH 8.5 ethanolamine chloride. The HLA/peptide complexes formed as indicated in paragraph I-1 are injected (30 µl) at a rate of 2 µl/min and will be able to bind to the immobilized antibody. The results are expressed in resonance units (RU) which correspond to the angle of deflection of the light beam, this deflection being proportional to the concentration of proteins in contact with the matrix. The system can be reused after regeneration by injecting 1 M ethanolamine for 2 min. Kinetic and affinity measurements can be carried out.

2) **Induction of cytolytic T lymphocytes *in vitro***

Unpurified peripheral blood mononuclear cells (PBMC) are cultured in RPMI 1640 and 10% BSA with antibiotics, at a concentration of 2×10^6 /ml, at a rate of 2 ml per well (Costar 24-well plates). 1 µg/ml tetanus toxin (TT) is added to these cells to stimulate the CD4+ T cells, and the peptide to be tested is added as a CTL inducer at a rate of 1 µg/ml. On day 3 or 4, 50 U/ml of IL-7 are added. For each culture, 10 replicates are treated independently. The effector cells generated are restimulated with PBMCs preincubated with the peptide (50 µg/ml for 4 h for 10×10^6 cells) and then irradiated at 6000 rads. The stimulator cells are diluted to 10^6 /ml and 1 ml is added per culture well after 1 ml of supernatant has been removed. The following day and 4 days later, IL-2 (10 U/ml) and IL-7 (50 U/ml) are added. The effectors are restimulated every 7 to 10 days in the same way. When the cytolytic activity appears, IL-2 is added at a rate of 50 U/ml. The cytotoxicity can be tested after 2 stimulations and then each week.

3) Analysis of the effector functions

3.1. Test of direct cytotoxicity

The induced T lines are tested as regards their capacity to recognize the inducer peptide presented on the HLA at the surface of the target cells, which are usually EBV lymphoblastoid cells. The cytotoxicity resulting therefrom is determined by the standard test of 4 h of release of ^{51}Cr . On day -3 or -4 before the test, the addition of interleukins is not carried out since the cells need to be weaned in order to be in optimum activation condition. For the primary inductions, the lysis cannot be detected before 3 weeks. The target cells labelled for 1 h with 10 μCi of ^{51}Cr are preincubated with the peptide (5 $\mu\text{g/ml}$ conventionally or a variable concentration). After 2 washes, these cells are distributed in the wells of a 96-well microtitration plate at a rate of 5000 per well and the effector cells are added in ratios ranging from 1 to 100. The release of ^{51}Cr (R) obtained during the incubation of 4 h at 37°C is measured. The percentage of lysis is determined by the formula below:

$$(\text{R. experimental} - \text{R. spontaneous} / \text{R. total} - \text{R. spontaneous}) \times 100$$

3.2. Analysis of the interleukins

Analysis of the expression of the genes coding for the interleukins is carried out by means of a semi-quantitative RT-PCR technique. PBMCs or purified CD8+ cells (1 to 2 $\times 10^6$) are cultured in 24-well plates and stimulated either with anti-CD3+ monoclonal antibodies in the presence of 100 nM Phorbol Myristate Acetate (PMA) with autologous PBMCs which have been irradiated and sensitized or otherwise for 1 h 30 min with a peptide whose recognition is restricted by the molecules of the MHC. For the various interleukins, after 12 to 14 h of stimulation (or after 3, 10 and 24 h for the kinetic analysis of the appearance of the mRNAs), the cells are recovered and the total RNA is extracted by the technique with RNazol (Bioprobe, Montreuil, France). The RNA (1 to 2 mg) is retrotranscribed with superscript II (Gibco BRL). The cDNAs synthesized are diluted to 1/5 with water and their concentration is quantified by competitive PCR with a plasmid (pQB.2) containing the sequence of B actin (237-bp product, generously supplied by D. Shire). The PCR reaction is carried out in the presence of 10 mM Tris-HCl, 50 mM KCl, 1.5 mM MgCl_2 , 0.2 mM of each dNTP (dATP, dCTP, dGTP, dTTP), 0.4 mM of sense and antisense primer, and 2 U of Taq polymerase (Promega). The amplification is carried out starting with a denaturation at 94°C for 5 min, followed by 30 cycles at 94°C, 30 s, 55°C, 30 s, 72°C, 30 s, and a final elongation at 72°C, 10 min, in a regulated thermostat (Perkin Elmers 9600). The analysis of IL-2, IL-4 and IFN- γ is carried out under the same conditions as for B-actin, except that the number

of cycles ranges between 30 and 40. The relative quantification is carried out by competitive PCR.

B) Results

5 1) Peptide analogues of the peptide M58-66 containing non-protein-generating amino acids :

1.1. Binding to the A2 molecule of the peptides of the series M58-66 :

The results are represented in Figure 1 (Figures 1A to 1C).

10 The peptides were incubated at different concentrations at 4°C for 18 h in the presence of denatured HLA-A2 molecules and β 2 microglobulin. The stable HLA-A2/peptide complexes formed at 4°C are captured with the antibody BB7.2 adsorbed onto a plate and are revealed with a second anti β 2 microglobulin antibody coupled to alkaline phosphatase. The quantity of HLA-A2/peptide
15 complexes formed is proportional to the intensity of the fluorescence detected with cytofluor. NP 383-391 is a negative control peptide, 0, peptide corresponds to the background noise in the absence of peptide. F.U. : arbitrary fluorescence unit.

Only two modifications affect the binding of these modified peptides to the molecule HLA-A2 : modification by the D-Phenyl in position 62 ([D-Phe62]M58-
20 66) and in position 64 ([D-Phe64]M58-66).

1.2. Capacity of these peptides to induce lysis of the target T2 cells by means of a line of cytotoxic T cells (CTL) which is specific for M58-66 and restricted with the molecule A-A2.1.

25 The results are represented in Figure 2 (Figures 2A to 2D).

The target T2 cells A2 were treated with chromium (Cr51) and then incubated for 2 h with 1 μ g/ml of peptide. After washing, the target cells (5000 per well) were incubated for 4 h with the CTL line at a ratio of 10 effector cells (E)/1 target cell (C). The specific lysis is expressed as a percentage of lysis as a function
30 of the concentration of peptide incubated with the targets.

The modifications Phg-62 and 64, DPhe-62 and 64, Cha62 and 64, Tic 62 and 64 have an effect on the recognition of the cytolytic T lymphocytes which are specific for the peptide M58-66.

35 1.3. Dose response.

The results are represented in Figure 3 (Figures 3A and 3B).

The target T2 cells A2 were treated with (Cr51) and then incubated for 1 h with various concentrations of peptide. After washing, the target cells (5000 per well) were incubated for 4 h with the CTL line at a ratio 10E/1C. The specific

lysis is expressed as a percentage of lysis as a function of the concentration of peptide incubated with the targets.

The cytolytic T lymphocytes which are specific for the peptide M58-66 have a weaker affinity for the modified peptides.

2) Peptide analogues of Mart-1 comprising a bond of the β -homologation type :

2.1. Binding to the molecule A2.

The results are represented in Figure 4 (Figures 4A and 4B).

The various peptides are compared as regards their ability to bind to the purified HLA-A2.1 molecule.

The peptides were incubated at different concentrations at 4°C for 18 h in the presence of denatured HLA-A2 molecules and β 2 microglobulin. The stable complexes HLA-A2/peptide formed at 4°C are captured with the antibody BB7.2 adsorbed onto a plate and revealed with a second anti β 2 microglobulin antibody coupled with alkaline phosphatase. The quantity of HLA-A2/peptide complexes formed is proportional to the intensity of the fluorescence detected with cytofluor (F.U.: arbitrary fluorescence unit).

NP is an influenza virus nucleoprotein peptide which does not bind to the molecule HLA-A2 (negative control).

Apart from the β 4 homo peptide which binds better than the parent peptide, the other β homo peptides have an intermediate to weak affinity relative to the peptide Mart-1.

2.2. Ability of these peptides to induce lysis of the target T2 cells with 3 T clones specific for Mart-1 which are restricted with the molecule HLA-A2.1.

The results are represented in Figure 5.

The target T2 cells A2 were treated with chromium (Cr51) and then incubated for 2 h with 1 μ g/ml of peptide. After washing, the targets (1500 per well) were incubated for 4 h with the various T clones (LT8, LT11, LT12) at a ratio of 3 effector cells/1 target cell (E/T). The specific lysis is expressed as a percentage of lysis.

The target T2 cells A2 incubated with the parent peptide Mart-1 are 90% lysed with the three cytolytic clones (LT8, LT11 and LT12) which are specific for this peptide. In the absence of peptide (0), no lysis is detected. Only the β 4 homo peptide is capable of inducing a lysis activity similar to that of the parent peptide for only one clone (LT12).

2.3. Dose response.

The results are represented in Figure 6.

The target T2 cells A2 were treated with chromium (Cr51) and then incubated for 2 h with various concentrations of peptide. After washing, the targets (1500 per well) were incubated for 4 h with the T clone LT12 at a ratio 3E/1C. The specific lysis is expressed as a percentage of lysis as a function of the concentration of peptide incubated with the targets.

The T clone LT12 has a weaker affinity for the β 4 peptide than for the peptide Mart-1.

3) Peptide analogues of the peptide Mart-1 containing a -CH₂-NH- bond.

3.1. Binding of the peptides of the reduced Mart-1 series to the molecule A2.

The results are represented in Figure 7 (Figures 7A and 7B).

Peptide/HLA-A2 complexes formed as a function of the concentration of peptide :

The peptides were incubated at two different concentrations (10^{-4} M and 10^{-6} M) at 4°C for 18 h in the presence of denatured HLA-A2 molecules and β 2 microglobulin. The stable HLA-A2/peptide complexes formed at 4°C are captured with the antibody BB7.2 (which is specific for this molecule of category I and is adsorbed at the bottom of the wells) and revealed by a second anti β 2 microglobulin antibody coupled with alkaline phosphatase. The quantity of HLA-A2/peptide complexes formed is proportional to the intensity of the fluorescence detected with cytofluor (F.U. : arbitrary fluorescence unit).

The peptides ψ 1-2 to 4-5 have an intermediate to weak affinity for the peptide Mart-1. The peptides ψ 4-5 to 8-9 bind very weakly to the purified A2.1 molecule.

3.2. Ability of the peptides of the reduced Mart series to induce lysis of the target T2 cells with 3 T clones specific for Mart which are restricted with the molecule HLA-A2.1.

The results are represented in Figure 8.

Lysis test :

Target T2 cells A2 were treated with chromium (Cr51) and then incubated for 2 h with 1 μ g/ml of peptide. After washing, the targets (1500 per well) were incubated for 4 h with the various T clones (LT8, LT11, LT12) at a ratio of 3 effector cells/1 target cell (E/T). The specific lysis is expressed as a percentage of lysis as a function of the various clones tested.

The target T2 cells incubated with the parent peptide Mart-1 are 90% lysed with the three cytolytic clones (LT8, LT11 and LT12) which are specific for this peptide. In the absence of peptide (0), no lysis is detected. Only the peptides ψ 2-3 and ψ 5-6 are capable of inducing a lysis activity similar to that of the parent peptide for a single clone (LT8 and LT12 respectively). ψ 7-8 induces a weaker lysis activity, detected only with the clone LT12.

3.3. Dose response of the peptides which induce a lysis with the clones LT8 and LT12.

The results are represented in Figure 9 (Figures 9A and 9B).

Lysis test :

Target T2 cells A2 were treated with chromium (Cr51) and then incubated for 2 h with various concentrations of peptide. After washing, the targets (1500 per well) were incubated for 4 h with the various T clones (LT8, LT12) at a ratio of 3E/1C. The specific lysis is expressed as a percentage of lysis as a function of the concentration of peptide incubated with the targets.

The peptides ψ 2-3 and ψ 5-6 have a dose response which is comparable to that of the parent peptide, whereas the peptide ψ 7-8 only induces lysis weakly, even at a high concentration of peptide. This result suggests the modification introduced reduces the affinity of the TCR (T receptor) of the clone LT8 for this peptide.

4) Peptide analogues of the peptide M58-66 containing a -CH₂-NH- bond.

4.1. Binding of the peptides of the reduced M58-66 series to the molecule A2.

The results are represented in Figure 10 (Figures 10A and 10B).

The peptides were incubated at various concentrations at 4°C for 18 h in the presence of denatured HLA-A2 molecules and β 2 microglobulin. The stable HLA-A2/peptide complexes formed at 4°C are captured with the antibody BB7.2 adsorbed on the plate and are revealed with a second anti β 2 microglobulin antibody coupled with alkaline phosphatase. The quantity of HLA-A2/peptide complexes formed is proportional to the intensity of the fluorescence detected with cytofluor. F.U. : arbitrary fluorescence unit.

The peptides Mart-1 modified by reduction have an intermediate affinity for the molecule HLA-A2. The reduction in positions 2-3 and 8-9 makes the binding undetectable for the clone studied.

4.2. Ability of the peptides of the reduced M58-66 series to induce lysis of the target T2 cells with a CTL line induced with M58-66 and restricted with the molecule HLA-A2.1.

The results are represented in Figure 11.

The target T2 cells A2 were treated with chromium (Cr51) and then incubated for 1 h with 1 μ g/ml of peptide. After washing, the targets (1500 per cell) were incubated for 4 h with the CTL line at a ratio of 10E/1C. The specific lysis is represented as a percentage of lysis.

The reduction in positions 2-3 and 5-6 does not have an effect on the recognition of the peptide by the cytolytic T lymphocytes which are specific for the peptide M58-66.

KEY TO THE FIGURES :

- Figure 1 : study of the binding of the peptide analogues of the peptide M58-66 containing non-protein-generating amino acids to the molecules of the MHC (HLA-A2) : the concentrations of the peptide analogues (10^{-4} to 10^{-10} M) are indicated on the x-axis and the fluorescence units are indicated on the y-axis :

. Figure 1A : study of the binding of the peptide analogues [Tic62]M58-66 (also known as M58-66 Tic62 or Tic62), [3-Pya62]M58-66 (3-Pya62), [2-Tha62]M58-66 (2-Tha62), [D-Phe62]M58-66 (D-Phe62), [Tic64]M58-66 (Tic64),

. Figure 1B : study of the binding of the peptide analogues [3-Pya64]M58-66 (3-Pya64), [2-Tha64]M58-66 (2-Tha64), [D-Phe64]M58-66 (D-Phe64), [Cha62]M58-66 (Cha62), [N-MePhe62]M58-66 (N-MePhe62),

. Figure 1C : binding of the peptide analogues [pCl-Ph62]M58-66 (pCl-Phe62), [Phg62]M58-66 (Phg62), [Cha64]M58-66 (Cha64), [pCl-Phe64]M58-66 (pCl-Phe64), [Phg64]M58-66 (Phg64),

in comparison with the peptide M58-66, the negative control peptide NP383-391, and a solution containing no peptide (0 peptide).

- Figure 2 : study of the effect of peptide analogues of M58-66 containing non-protein-generating amino acids on the lysis of a line of target T2 cells with a line of cytotoxic T cells which are specific for M58-66 : the percentage of lysis is indicated on the y-axis and the ratio E/T (effector cells/target cells) is represented on the x-axis:

Figure 2A : effect of the analogues Tic62 and Tic64 on the lysis of the T2 cells,

Figure 2B : effect of the analogues D-Phe62, D-Phe64, Cha62 and Cha64 on the lysis of the T2 cells,

. Figure 2C : effect of the analogues 3-Pya62, 3-Pya64, 2-Tha62 and 2-Tha64 on the lysis of the T2 cells,

. Figure 2D : effect of the analogues pCl-Phe64, Phg64, pCl-Phe62 and Phg62 on the lysis of the T2 cells,
in comparison with the effects of the peptide M58-66 and of a solution containing no peptide (0 peptide).

- Figure 3 : study of the response of cytotoxic T cells as a function of the concentration of peptide analogues of the peptide M58-66 containing non-protein-generating amino acids : the percentage of lysis is indicated on the y-axis and the peptide concentration is indicated on the x-axis :

. Figure 3A : study as a function of the concentration of pCl-Phe62 and pCl-Phe64,

. Figure 3B : study as a function of the concentration of 2-Tha62 and 2-Tha64,

in comparison with the concentration of M58-66.

- Figure 4 : study of the binding of the peptide analogues of the peptide Mart-1 containing a bond of the β -homologation type to the molecules of the MHC (HLA-A2); the concentrations of the peptide analogues (10^{-5} to 10^{-7} M) are indicated on the x-axis and the fluorescence units are indicated on the y-axis :

. Figure 4A : study of the binding of the analogues $\beta 1$ homo (B1), $\beta 2$ homo (B2), $\beta 3$ homo (B3) and $\beta 4$ homo (B4),

. Figure 4B : study of the binding of the analogues $\beta 5$ homo (B5), $\beta 6$ homo (B6), $\beta 7$ homo (B7), $\beta 8$ homo (B8) and $\beta 9$ homo (B9),

in comparison with the peptide Mart-1 (Mart) and a negative control peptide (NP).

. Figure 5 : study of the effect of peptide analogues of the peptide Mart-1 containing a bond of the β -homologation type on the lysis of target T2 cells with 3 clones of cytotoxic T cells (LT8, LT11 and LT12) which are specific for Mart-1: the percentage of lysis is indicated on the y-axis and the peptide analogues B1 to B9 are indicated on the x-axis, in comparison with Mart-1 and a solution without peptide (0).

- Figure 6 : study of the response of the cytotoxic T cells as a function of the concentration of the $\beta 4$ homo analogue of Mart-1, in comparison with Mart-1 : the percentage of lysis is indicated on the y-axis and the concentration of peptide ($\mu\text{g/ml}$) is indicated on the x-axis.

- Figure 7 : study of the binding of the peptide analogues of the peptide Mart-1 containing a $-\text{CH}_2\text{-NH}-$ bond (reduced analogues) to the molecules of the MHC (HLA-A2) : the peptides at concentrations of 10^{-6} M or 10^{-4} M are indicated on the x-axis and the fluorescence units are indicated on the y-axis :

. Figure 7A : study of the binding of the peptides $\psi(1-2)\text{Mart-1}$ (represented by 1-2), $\psi(2-3)\text{Mart-1}$ (2-3), $\psi(3-4)\text{Mart-1}$ (3-4) and $\psi(4-5)\text{Mart-1}$ (4-5),

. Figure 7B : study of the binding of the peptides $\psi(5-6)$ Mart-1 (5-6), $\psi(6-7)$ Mart-1 (6-7), $\psi(7-8)$ Mart-1 (7-8) and $\psi(8-9)$ Mart-1 (8-9), in comparaisn with Mart-1.

5 - Figure 8 : study of the effect of the reduced analogues 1-2 to 8-9 of Mart-1 on the lysis of T2 cells with 3 clones of T cells (LT8, LT11 and LT12) which are specific for Mart-1 : the percentage of lysis is indicated on the y-axis and the peptide analogues 1-2 to 8-9 are indicated on the x-axis, in comparison with Mart-1 and a solution without peptide (0).

10 - Figure 9 : study of the response of the cytotoxic T cells as a function of the concentration of reduced analogues of Mart-1: the percentage of lysis is indicated on the y-axis and the peptide concentration is indicated on the x-axis ($\mu\text{g/ml}$) :

. Figure 9A : study as a function of the concentration of the analogues 2-3 and 7-8,

15 . Figure 9B : : study as a function of the concentration of the analogue 5-6, in comparison with the concentration of Mart-1.

- Figure 10 : study of the binding of peptide analogues of the peptide M58-66 containing a $-\text{CH}_2\text{-NH-}$ bond (reduced analogues) to the molecules of the MHC HLA-A2) : the peptide concentrations (10^{-10} to 10^{-4} M) are indicated on the x-axis and the fluorescence units are indicated on the y-axis :

20 . Figure 10A : study of the binding of the peptides $\psi(1-2)$ M58-66 (1-2), $\psi(2-3)$ M58-66 (2-3), $\psi(3-4)$ M58-66 (3-4) and $\psi(4-5)$ M58-66 (4-5),

. Figure 10B : study of the binding of the peptides $\psi(5-6)$ M58-66 (5-6), $\psi(6-7)$ M58-66 (6-7), $\psi(7-8)$ M58-66 (7-8) and $\psi(8-9)$ M58-66 (8-9),

25 in comparisn with M58-66 and a mutated peptide (GIL) of M58-66 bearing the mutation $\text{GLL} \rightarrow \text{GIL}$ [M58-66 (GIL)],

30 - Figure 11 : study of the effect of the reduced analogues 1-2 to 8-9 of M58-66 on the lysis of T2 cells with a clone of T cells which is specific for M58-66 : the percentage of lysis is indicated on the y-axis and the effector cell/target cell (E/T) ratio is indicated on the x-axis, in comparison with M58-66, mutated (GLL) M58-66 and a solution without peptide (0).

BIBLIOGRAPHIC REFERENCES

- 5
1. Bohnstedt A.C. *et al.* Tetrahedron Lett. 1993, 34:5217-5220.
2. Breton P. *et al.* Int. J. Peptide Protein Res. 1990, 35:346-351.
3. Chorev M. & Goodman M. Acc. Chem.. Res. 1993, 26:266-273.
- 10
4. DeMars *et al.* Proc. Natl. Acad. Sci. USA 1985, 82:8183.
5. Evans B.E. *et al.* J. Org. Chem. 1985, 50:4615-4625.
6. Gnjatic S. *et al.* Eur. J. Immunol. 1995, 25:1638-1642.
- 15
7. Gonzalez-Muniz R. *et al.* J. Med. Chem. 1995, 38:1015-1021.
8. Goodman M. & Chorev M. Acc. Chem. Res. 1979, 12:1-7.
- 20
9. Harbeson S.L. & Rich D.H. J. Med. Chem. 1989, 32:1378-1392.
10. Hann M.M. *et al.* J. Chem. Soc. Perkin Trans. 1982, 1:307-314.
11. Kempf D.J. *et al.* Int. J. Peptide Protein Res. 1991, 38:237-241.
- 25
12. Ljunggren H.G. & Kärre K. J. Exp. Med. 1985, 162:1745.
13. Martinez J. *et al.* J. Med. Chem. 1985, 28:1874.
- 30
14. Parham P. & Brodsky F.M. J. Immunol. 1981, 3:277.
15. Rodriguez M. *et al.* Tetrahedron Lett. 1990a, 31:5153-5156.
16. Rodriguez M. *et al.* Tetrahedron Lett. 1990b, 31:7319-7322.
- 35
17. Salter R.D. & Cresswell P. EMBO J. 1986, 5:943.
18. Sasaki Y. & Coy D.H. Peptides 1987, 8:119-121.

19. Shemyakin M.M. *et al.* Angew Chem. Internat. Edit. 1969, 8:492-499.
20. Szelke M. *et al.* Nature 1982, 299:555-557.
21. Teillaud *et al.* Immunogenetics 1982, 15:377.
22. TenBrink R.E. J. Org. Chem. 1987, 52:418-422.
23. Wuts P.G.M. *et al.* J. Org. Chem. 1992, 57:6696-6700.

5

10

[illegible]

CLAIMS

1. Use of peptide analogues of parent peptides, these parent peptides being derived, where applicable, from exogenous or endogenous proteins, the said parent peptides interacting with molecules of the MHC in the context of pathologies involving an immune response to cell mediation, in man or animals, the said analogues being characterized in that they correspond to the said parent peptides in which:

- at least one peptide bond -CO-NH- of the peptide chain is modified, with the exception of modifications of the retro or retro-inverso type, or

- at least one amino acid of the peptide chain is substituted with a non-protein-generating amino acid, or



- at least one peptide bond -CO-NH- of the peptide chain is modified and at least one amino acid of the said peptide chain is substituted with a non-protein-generating amino acid,

for the preparation of a medicinal product intended for preventing or treating the abovementioned pathologies.

2. Use according to Claim 1, of peptide analogues derived from parent peptides which interact with molecules of the MHC of category I, in the context of pathologies involving cytotoxic T lymphocytes.

3. Use according to Claim 1 or 2, of peptide analogues, characterized in that at least one of the peptide bonds -CO-NH- in the peptide chain of the parent peptide is replaced with a bond other than the -CO-NH-, the said other bond being chosen in particular from the following :

- | | |
|-------------------------------------|-------------------------|
| -CH ₂ -NH- | (methylenamino) ; |
| -CH ₂ -CH ₂ - | (carba) ; |
| -CO-CH ₂ - | (ketomethylene) ; |
| -CH ₂ -O- | (methyleneoxy) ; |
| -CHOH-CH ₂ - | (hydroxyethylene) ; |
| -CH=CH- | (E or Z olefin) ; |
| -CHOH-CHOH- | (dihydroxyethylene) ; |
| -CHCN-NH- | (cyanomethyleneamino) ; |
| -S-CH ₂ - | (thiomethylene) ; |
| -CH ₂ -S- | (methylenethio) ; |
| -CS-NH- | (thioamide) ; |
| -PO ₂ -NH- | (phosphonamide) ; |

- CHOH- (hydroxymethylene) ;
 -NH-CO-NH- (urea) ;
 -CH-CH- (oxirane) ;

 5 -C-N- (tetrazole) ;

 -CH₂-CO-NH- (β-homologation) ;
 -CHOH-CH₂-NH- (hydroxyethyleneamino) ;
 10 -CO-NH-NH- (hydrazino).

4. Use according to one of Claims 1 to 3, of peptide analogues, characterized in that at least one of the amino acids in the peptide chain of the parent peptide is substituted with a non-protein-generating amino acid, the said non-protein-generating amino acid being chosen in particular from the following amino acids:

- the amino acids of D configuration,
- the α-amino acids of the general formula:


in which :

- R₁, R₂ and R₃, represent, independently of each other : a hydrogen atom, a hydroxyle, an alkyl radical of 1 to 25 carbon atoms, a radical containing an allyl group and having from 3 to 25 carbon atoms, a radical containing one or more aromatic or non-aromatic rings, in particular an aryl group, and having from 6 to 25 carbon atoms, and in particular the following groups: -CH₃ (methyl), -CH₂CH₃ (ethyl), -(CH₂)₄-CH₃, -CH(CH₃)₂ (isopropyl), -C(CH₃)₃ (tert-butyl), -Φ (phenyl), -CH₂ Φ (benzyl), -CH₂ ΦCl (para-chlorobenzyl), -CH₂-CH₂ Φ (2-phenylethyl), -CH₂CHCH₂ (alkyl), methylfluorenyl, -CH₂CONH₂ (glycolamide), -CH₂CON Φ₂ (benzhydrylglycolamide), -CHOH Φ,




it being understood that one of the two groups R₂ and R₃ can represent a side chain of natural amino acids when either R₁ or the other of the two groups R₂ and R₃ do not represent a hydrogen atom,

- where appropriate, R_1 , R_2 , $C\alpha$ and N form an aromatic or non-aromatic heterocycle of 4 to 8 carbon atoms, which may be substituted, in particular a heterocycle of the formula :




10 - the β -amino acids of the general formula :



15 in which R_1 , R_2 and R_3 , independently of each other, represent a side chain of a natural amino acid or are as defined above,


in particular :

the β -homo amino acids of the formula:




in which R_1 and R_2 are as defined above, or

the α -hydroxy β -homo amino acids of the formula :




in which R_1 and R_2 are as defined above,

30 - the γ -amino acids of the general formula:



35 in which R_1 , R_2 , R_3 and R_4 represent, independently of each other, a side chain of a natural amino acid, or R_1 , R_2 and R_3 , are as defined above and R_4 has the same meaning as that given above for R_1 , R_2 and R_3 ,

in particular the statin derivatives of the formula :



5

5. Use according to one of Claims 1 to 4, of peptide analogues, characterized in that they are selected from those which are capable :

- on the one hand, of being recognized by the molecules of the MHC and of associating with these molecule, in particular by carrying out the following method :

10

. incubation of the peptide analogue in the presence of molecules of the MHC, derived from the lysis of human or animal cells, or purified in particular by affinity chromatography from human or animal cell lines, on a solid support coated with a first antibody, in particular a monoclonal antibody, which specifically recognizes the molecules of the MHC in their conformation which is dependent on their binding to the said peptide analogue

15

. addition to the above solid support of a second antibody which is labelled, in particular by means of coupling with a radioactive, enzymatic or fluorescent label, the said labelled antibody specifically recognizing either the molecules of the MHC in their conformation which is dependent on their binding to the peptide analogue, or a molecule which itself binds specifically to the molecules of the MHC in their abovementioned conformation, in particular β 2-microglobulin which specifically recognizes the molecules of the MHC of category I,

20

. detection, after rinsing the solid support, of the possible presence of the second labelled antibody which has remained bound to the solid support, thereby demonstrating an effect of recognition and association between the molecules of the MHC and the peptide analogue studied,

25

- and, on the other hand, of forming a complex with the said molecules of the MHC, the stability of which complex can be evaluated by carrying out a method for monitoring over time the binding established between the peptide analogue and the molecules of the MHC, this method advantageously being carrying out according to a protocol which is identical to the above method, but in which the step of incubation of the peptide analogue in the presence of the molecules of the MHC on

30

007550 2336460

the solid support coated with the said first antibody is carried out for times ranging from a few minutes to several days.

5 6. Use according to one of Claims 1 to 5, of peptide analogues, characterized in that they are selected :

* from those which are capable :

- of inducing *in vitro* the appearance and growth of cytotoxic T lymphocytes from human or animal cells, in particular from peripheral blood mononuclear cells (PBMCs) in the presence of factors required for the growth and
10 differentiation of the cytotoxic T cells,

- of inducing *in vitro* cytotoxicity, by means of cytotoxic T lymphocytes, of target cells carrying at their surface the peptide analogue associated with the molecules of the MHC, the said cytotoxic T lymphocytes advantageously being taken from a patient suffering from a pathology in which the parent peptide of the
15 peptide analogue studied is involved,

- and of inducing *in vitro* the secretion of cytokines (or interleukins) by means of the abovementioned cytotoxic T lymphocytes, in particular IL-2, IL-4 or γ -interferon,

the said peptide analogues thus selected being :

20 . either receptor agonists (TCR) which recognize the antigen (i.e. the parent peptide) of the cytotoxic T cells, and are derived from parent peptides which themselves behave as agonists or antagonists of the said receptors,

. or partial agonists of the said receptors, and are derived from parent peptides which themselves behave as agonists of the said receptors, these partial
25 agonists inducing, in particular, the secretion of one or more cytokines other than those whose secretion is induced with the parent peptides,

* or from those :

- which are capable of inducing *in vitro* the appearance and growth of cytotoxic T lymphocytes from human or animal cells, in particular from peripheral
30 blood mononuclear cells (PBMCs), in the presence of factors required for the growth and differentiation of cytotoxic T cells,

- which do not induce *in vitro* the cytotoxicity, by means of cytotoxic T lymphocytes, of target cells carrying at their surface the peptide analogue associated with the molecules of the MHC, the said cytotoxic T lymphocytes advantageously

0940357-05400

being taken from a patient suffering from a pathology in which the parent peptide of the peptide analogue studied is involved,

- which do not induce *in vitro* the secretion of cytokines (or interleukins) by means of the abovementioned cytotoxic T lymphocytes, in particular IL-2, IL-4 or γ -interferon,

the said peptide analogues thus selected being antagonists of the cytotoxic T cell receptors.

7. Peptide analogues of parent peptides involved in melanoma, in particular the peptide MART1 27-35, the said parent peptide comprising, where appropriate, one or more mutations, such as the Leu²⁸-mutated parent peptide MART1 27-35, the said peptide analogues corresponding to the said parents peptides in which at least one of the peptide bonds -CO-NH- is modified, with the exception of modifications of the retro or retro-inverso type, the said analogues being chosen in particular from the following :

- the peptide analogues of MART1 27-35 in which at least one of the -CO-NH- bonds is replaced with a -CH₂-NH- bond, such as the analogues Ψ (1-2) to Ψ (8-9) below :

		Sequences								
		P1	P2	P3	P4	P5	P6	P7	P8	P9
25	MART1 27-35	H - A -	A -	G -	I -	G -	I -	L -	T -	V - OH
	Ψ (1 - 2)	H - A	Ψ (CH ₂ NH)A -	G -	I -	G -	I -	L -	T -	V - OH
	Ψ (2 - 3)	H - A - A	Ψ (CH ₂ NH)G -	I -	G -	I -	L -	T -	V - OH	
	Ψ (3 - 4)	H - A -	A - G	Ψ (CH ₂ NH)I -	G -	I -	L -	T -	V - OH	
30	Ψ (4 - 5)	H - A -	A	G -	I	Ψ (CH ₂ NH)G -	I -	L -	T -	V - OH
	Ψ (5 - 6)	H - A -	A	G -	I -	G	Ψ (CH ₂ NH)I -	L -	T -	V - OH
	Ψ (6 - 7)	H - A -	A	G -	I -	G	I	Ψ (CH ₂ NH)L -	T -	V - OH
	Ψ (7 - 8)	H - A -	A	G -	I -	G	I -	L	Ψ (CH ₂ NH)T -	V - OH
	Ψ (8 - 9)	H - A -	A	G -	I -	G	I -	L -	T	Ψ (CH ₂ NH)V - OH
35										

- the peptide analogues of MART1 27-35 in which at least one of the -CO-NH- bonds is replaced with a -CH₂-CO-NH- bond, also referred to hereinbelow as a β -homo bond, such as the analogues β 1 to β 9 below :

40


Sequences

	P1	P2	P3	P4	P5	P6	P7	P8	P9
MARTI 27-35	H-A	A-	G-	I-	G-	I-	L-	T-	V-OH
5 β 1	H- β -homoA	-A-	G-	I-	G-	I-	L-	T-	V-OH
β 2	H-A	- β -homoA-	G-	I-	G-	I-	L-	T-	V-OH
β 3	H-A-	A-	β -homoG-	I-	G-	I-	L-	T-	V-OH
β 4	H-A-	A-	G-	β -homoI-	G-	I-	L-	T-	V-OH
β 5	H-A-	A-	G-	I-	β -homoG-	I-	L-	T-	V-OH
10 β 6	H-A-	A-	G-	I-	G-	β -homoI-	L-	T-	V-OH
β 7	H-A-	A-	G-	I-	G-	I-	β -homoL-	T-	V-OH
β 8	H-A-	A-	G-	I-	G-	I-	L-	β -homoT-	V-OH
β 9	H-A-	A-	G-	I-	G-	I-	L-	T-	β -homoV-OH


15 - the peptide analogues of MART1 27-35 Leu²⁸ in which at least one of the -CO-NH- bonds is replaced with a -CH₂-CO-NH- bond, such as the abovementioned analogues β 1 to β 9 in which the alanine in P2 is replaced with a leucine,

- the peptide analogues of MART1 27-35 in which at least one of the -CO-NH- bonds is replaced with a -CH₂-CH₂- bond, such as the following analogues :


20



25








30



35

40











8. Peptide analogues of the parent peptides of the influenza virus, in particular of the parent peptide M58-66, the said peptide analogues corresponding to the said parent peptides in which:

- at least one of the -CO-NH- peptide bonds is modified, with the exception of modifications of the retro or retro-inverso type, the said analogues being chosen in particular from those in which at least one of the -CO-NH- bonds is replaced with a -CH₂-NH- bond, such as the following analogues :


Sequences	
	P1 P2 P3 P4 P5 P6 P7 P8 P9
M58-66	H- G - L - L - G - F - V - F - T - L -OH
Ψ(1-2)	H-GΨ(CH ₂ NH)L-L- G - F - V - F - T - L -OH
Ψ(2-3)	H- G- LΨ(CH ₂ NH)L- G - F - V - F - T - L -OH
35 Ψ(3-4)	H- G - L-LΨ(CH ₂ NH)G- F - V - F - T - L -OH
Ψ(4-5)	H- G - L - L - GΨ(CH ₂ NH)F- V - F - T - L -OH
Ψ(5-6)	H- G - L - L - G - FΨ(CH ₂ NH)V- F - T - L -OH
Ψ(6-7)	H- G - L - L - G - F - VΨ(CH ₂ NH)F- T - L -OH
Ψ(7-8)	H- G - L - L - G - F - V - FΨ(CH ₂ NH)T - L -OH
40 Ψ(8-9)	H- G - L - L - G - F - V - F - TΨ(CH ₂ NH)L-OH

- at least one of the amino acids of the peptide chain is substituted with a non-protein-generating amino acid, such as the following analogues :




61


[2-Tha64]M58-66




[D-Phe64]M58-66




[Cha62]M58-66




[N-MePhe62]M58-66




[pCl-Phe62]M58-66




[Phg62]M58-66




[Cha64]M58-66



[pCl-Phe64]M58-66



[Phg64]M58-66



9. Peptide analogues of parent peptides of the AIDS virus, in particular of the peptides NEF 84-92 and GAG 77-85, the said peptide analogues corresponding to the said parent peptides in which at least one of the -CO-NH- peptide bonds is modified, with the exception of modifications of the retro or retro-inverso type, the said analogues being chosen in particular from :

- those in which at least one of the -CO-NH- bonds is replaced with a -CH₂-NH-, such as the analogues NEFRD1-8 of the peptide NEF, and GAGRD1-8 of the peptide GAG, below:

NEF	AVDLSHFLK
NEFRD1	AΨ(CH ₂ -NH)VDLSHFLK
NEFRD2	AVΨ(CH ₂ -NH)DLSHFLK
NEFRD3	AVDΨ(CH ₂ -NH)LSHFLK
NEFRD4	AVDLΨ(CH ₂ -NH)SHFLK
NEFRD5	AVDLSΨ(CH ₂ -NH)HFLK
NEFRD6	AVDLSHΨ(CH ₂ -NH)FLK
NEFRD7	AVDLSHFΨ(CH ₂ -NH)LK
NEFRD8	AVDLSHFLΨ(CH ₂ -NH)K

	GAG	SLYNTVATL
	GAGRD1	SΨ(CH₂-NH)LYNTVATL
	GAGRD2	SLΨ(CH₂-NH)YNTVATL
5	GAGRD3	SLYΨ(CH₂-NH)NTVATL
	GAGRD4	SLYNΨ(CH₂-NH)TVATL
	GAGRD5	SLYNTΨ(CH₂-NH)VATL
	GAGRD6	SLYNTVΨ(CH₂-NH)ATL
	GAGRD7	SLYNTVAΨ(CH₂-NH)TL
10	GAGRD8	SLYNTVATΨ(CH₂-NH)L

- those in which at least one of the -CO-NH- bonds is replaced with a -CHOH-NH- bond, such as the following analogues NEFHEA1-8 of the peptide NEF:

15	NEF	AVDLSHFLK
	NEFHEA1	AΨ(CHOH-NH)VDLSHFLK
	NEFHEA2	AVΨ(CHOH-NH)DLSHFLK
	NEFHEA3	AVDΨ(CHOH-NH)LSHFLK
	NEFHEA4	AVDLΨ(CHOH-NH)SHFLK
20	NEFHEA5	AVDLSΨ(CHOH-NH)HFLK
	NEFHEA6	AVDLSHΨ(CHOH-NH)FLK
	NEFHEA7	AVDLSHFΨ(CHOH-NH)LK
	NEFHEA8	AVDLSHFLΨ(CHOH-NH)K

25 **10.** Use of peptide analogues as defined in one of Claims 1 to 9, for the preparation of medicinal products, in particular vaccines, intended for preventing or treating pathologies in which the parent peptides are agonists or antagonists of receptors which recognize the antigen of the cytotoxic T cells, and more particularly neurodegenerative pathologies which are infectious (of viral or bacterial origin),
30 tumoural, autoimmune and allergic.

35 **11.** Pharmaceutical composition, characterized in that it comprises, as active principle, at least one peptide analogue defined in one of Claims 1 to 9, in combination with a pharmaceutically acceptable vehicle.

12. Vaccine, characterized in that it comprises an agonist peptide analogue, which may be a partial agonist, defined in one of Claims 1 to 9, in combination with a pharmaceutically acceptable vehicle.

5 13. Diagnostic composition for the *in vivo* diagnosis of pathologies involving the immune response to cell mediation, in particular the cytotoxic T lymphocytes, or for the *in vivo* evaluation of the immune response in the context of the abovementioned pathologies, by carrying out a skin hypersensitivity reaction by means of intradermal injection of the said diagnostic composition, characterized in
10 that it comprises a peptide analogue defined in one of Claims 1 to 9, in combination with a physiologically acceptable vehicle.

14. Complex between a peptide analogue defined in one of Claims 1 to 9, and a component of the major histocompatibility complex (also referred to as an MHC-peptide analogue binary complex), and optionally a T cell receptor (also referred to
15 as an MHC-peptide analogue-T receptor ternary complex).

15. Method for the *in vitro* diagnosis of pathologies involving the immune response to cell mediation, in particular the cytotoxic T lymphocytes, i.e. pathologies
20 associated with the presence, in a patient's body, of exogenous or endogenous peptides which interact with molecules of the MHC, and which are liable to be directly or indirectly involved in the process of development of these pathologies in man or animals, characterized in that it comprises:

- placing a biological sample, obtained from a patient, in contact with a peptide
25 analogue defined in one of Claims 1 to 9, under conditions which allow reaction between the receptors of the T cells liable to be present in the biological sample, and the binary complex which may be formed between the said peptide analogue and the molecules of the MHC present in the said sample;

- the *in vitro* detection of the MHC-peptide analogue-T receptor ternary
30 complex which may be formed in the preceding step.

16. Equipment or kit for carrying out the methods of *in vitro* diagnosis according to Claim 15, comprising:

- a peptide analogue defined in one of Claims 1 to 9;

- reagents for making a medium suitable for forming an immunological reaction;

- reagents for detecting the ternary complex according to Claim 14, which has been produced after the immunological reaction, the said reagents optionally containing a label or being capable of being recognized in turn by a labelled reagent, more particularly in the case in which the peptide analogue is not labelled.

17. Antibodies directed against the binary complexes as defined in Claim 14, the said antibodies being as obtained by immunizing an animal with at least one abovementioned binary complex, the said antibodies being capable of forming a complex with these binary complexes.

18. Pharmaceutical composition, characterized in that it comprises antibodies according to Claim 17, in combination with a physiologically acceptable vehicle.

19. Process for screening peptide analogues defined in one of Claims 1 to 9, characterized in that it comprises the following steps :

. incubation for times ranging from a few minutes to several days, of the peptide analogue in the presence of molecules of the MHC, derived from the lysis of human or animal cells, or purified in particular by affinity chromatography from human or animal cell lines, on a solid support coated with a first antibody, in particular a monoclonal antibody, which specifically recognizes the molecules of the MHC in their conformation which is dependent on their binding to the said peptide analogue,

. addition to the preceding solid support of a second antibody which is labelled, in particular by means of coupling with a radioactive, enzymatic or fluorescent label, the said labelled antibody specifically recognizing either the molecules of the MHC in their conformation which is dependent on their binding to the peptide analogue, or a molecule which itself binds specifically to the molecules of the MHC in their abovementioned conformation, in particular β 2-microglobulin which specifically recognizes the molecules of the MHC of category I,

. detection, after rinsing the solid support, of the possible presence of the second labelled antibody which has remained bound to the solid support,

. evaluation of the duration of the association between the said peptide analogue and the molecules of the MHC.

20. Set or kit for carrying out a screening process according to Claim 19, comprising:

- molecules of the MHC, and/or
- 5 - antibodies which specifically recognize the molecules of the MHC in their conformation which is dependent on their binding to the said peptide analogue, which antibodies are advantageously bound to a solid support, or are supplied with the reagents required for binding them to the solid support, and/or
- 10 - antibodies which are labelled, in particular by means of coupling with a radioactive, enzymatic or fluorescent label, this antibody specifically recognizing either the molecules of the MHC in their conformation which is dependent on their binding to the peptide analogue, or a molecule which itself binds specifically to the molecules of the MHC in their abovementioned conformation, in particular ###2-microglobulin which specifically recognizes the molecules of the MHC of category
- 15 I, and/or
- a protocol for carrying out the said process, and/or
- a control peptide.

20

007659 / 2350460

ABSTRACT

5 **PEPTIDE ANALOGUES, ANTIBODIES DIRECTED AGAINST THE
LATTER, AND USES THEREOF, IN PARTICULAR IN
PHARMACEUTICAL AND DIAGNOSTIC COMPOSITIONS**

The invention relates to the use of peptide analogues of parent peptides, characterized in that they correspond to the said parent peptides in which :



- 10 - at least one peptide bond -CO-NH- of the peptide chain is modified, with the exception of modifications of the retro or retro-inverso type, or
- at least one amino acid of the peptide chain is substituted with a non-protein-generating amino acid, or
- 15 - at least peptide bond -CO-NH- of the peptide chain is modified and at least one amino acid of the said peptide chain is substituted with a non-protein-generating amino acid,
- for
- the preparation of a medicinal product intend for preventing or treating the abovementioned pathologies, or
- 20 - the preparation of a composition intended for the *in vivo* diagnosis of the abovementioned pathologies, or for the *in vivo* evaluation of the immune response in the context of the abovementioned pathologies, by carrying out a skin hypersensitivity reaction by means of intradermal injection of the said composition,
- carrying out a method for the *in vitro* diagnosis of the said pathologies.

25

(No figures)


09/403627

1 / 15

FIG. 1AFIG. 1B

09/403627

2 / 15

FIG.1C





FIG.2A

09/403627

4 / 15




FIG. 3A

09/403627

5 / 15




FIG. 2C




FIG. 2D

09/403627

7 / 15



FIG. 3AFIG. 3B

FIG.4A





FIG.4B






FIG.5

9/15

09/403627




FIG. 6

10 / 15

09/403627




FIG. 7A




FIG. 7B

09/11/99 12:56 09/403627

12/15

09/403627






FIG. 8

09/403627

13 / 15

FIG.9A



LT12








FIG.9B

09/403627

14 / 15



REPLACEMENT SHEET/PAGE



Attorney's Docket No. 759-11**COMBINED DECLARATION AND POWER OF ATTORNEY**(ORIGINAL, DESIGN, NATIONAL STAGE OF PCT, SUPPLEMENTAL,
DIVISIONAL, CONTINUATION OR CIP)

As a below named inventor, I hereby declare that:

TYPE OF DECLARATIONThis declaration is of the following type: *(check one)*

☐ Original
☐ Supplemental
☐ Design

☒ National Stage PCT
☐ Divisional
☐ Continuation
☐ Continuation-in-Part (CIP)

INVENTORSHIP IDENTIFICATION

NOTE If the inventors are each not the inventors of all the claims an explanation of the facts, including the ownership of all the claims at the time the last claimed invention was made, should be submitted

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

**PEPTIDE ANALOGUES AND USES THEREOF, IN PARTICULAR IN
 PHARMACEUTICAL AND DIAGNOSTIC COMPOSITIONS**

the specification of which: *(complete (a), (b) or (c))*(a) ☐ is attached hereto.(b) ☒ was filed on November 5, 1999 as☒ Serial No 09/403,627 or☐ Express Mail No. _____, as Serial No. not yet known
and was amended on Nov. 5, '99 *(if applicable)*(c) ☒ was described and claimed in PCT International Application No. PCT/FR98/00923
filed on May 7, 98 and as amended under PCT Article 19 on _____. *(If any)*

ACKNOWLEDGMENT OF REVIEW OF PAPERS AND DUTY OF CANDOR

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above, and that the filing of said specification, if heretofore filed, was authorized by me.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

CLAIM OF PRIORITY OF EARLIER FOREIGN APPLICATION(S) UNDER 35 U.S.C. §119(a)-(d)

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

(List prior foreign/PCT application(s) filed within 12 months (6 months for design) prior to this U S application)

NOTE Where item (c) is entered above and the International Application which designated the U.S. claimed priority check item (e), enter the details below and make the priority claim.

COUNTRY (or PCT)	APPLICATION NO. (Day/Month/Year)	DATE OF FILING	PRIORITY CLAIMED UNDER 35 USC §119
France	97/05677	May 7, 1997	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO

CLAIM FOR BENEFIT OF PRIOR U.S. PROVISIONAL APPLICATION(S) UNDER 35 U.S.C. §119(e)

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below:

(List prior U S provisional applications.)

PROVISIONAL APPLICATION NO.	FILING DATE (Day/Month/Year)

CLAIM FOR BENEFIT OF EARLIER U.S./PCT APPLICATION(S) UNDER 35 U.S.C. 120

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in such prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

(List prior U.S. applications or PCT international applications designating the U.S. for benefit under 35 U.S.C. §120.)

U.S. APPLICATIONS

STATUS (Check One)

U.S. SERIAL NO.	U.S. FILING DATE (Day/Month/Year)	Patented	Pending	Abandoned
0 /		<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
0 /		<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

PCT APPLICATIONS DESIGNATING THE U.S.

STATUS (Check One)

PCT APPLN. NO.	PCT FILING DATE (Day/Month/Year)	U.S. SERIAL NOS. ASSIGNED (If any)	Patented	Pending	Abandoned
PCT/			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
PCT/			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

35 USC 119 PRIORITY CLAIM, IF ANY, FOR ABOVE LISTED U.S./PCT APPLICATIONS

PRIORITY APPLICATION NO.	PRIORITY COUNTRY	FILING DATE (Day/Month/Year)	ISSUE DATE (Day/Month/Year)

POWER OF ATTORNEY

As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office in connection therewith:

Charles R. Hoffmann, Reg. No. 24,102; Ronald J. Baron, Reg. No. 29,281; Gerald T. Bodner, Reg. No. 30,449; Alan M. Sack, Reg. No. 31,874; A. Thomas Kammer, Reg. No. 28,226; R. Glenn Schroeder, Reg. No. 34,720; Glenn T. Henneberger, Reg. No. 36,074; Irving N. Feit, Reg. No. 28,601; Anthony E. Bennett, Reg. No. 40,910; Gregory W. Bachmann, Reg. No. 41,593; Steven T. Zuschlag, Reg. No. 43,309; Susan A. Sipos, Reg. No. 43,128; Kevin E. McDermott, Reg. No. 35,946; Robert C. Morriss, Reg. No. 42,910; Roderick S.W. Turner, Reg. No. 38,639; James F. Harrington, Reg. No. 44,741; Samir R. Patel, Reg. No. 44,998; Richard LaCava, Reg. No. 41,135; Algis Anilionis, Reg. No. 36,995; and Justin K. Holmes, Reg. No. 42,666, each of them of HOFFMANN & BARON, LLP, 6900 Jericho Turnpike, Syosset, New York 11791; and Daniel A. Scola, Jr., Reg. No. 29,855; Salvatore J. Abbruzzese, Reg. No. 30,152; Kirk M. Miles, Reg. No. 37,891; Robert F. Chisholm, Reg. No. 39,939; Kellyanne Merkel, Reg. No. 43,800; Keith R. Lange, Reg. No. 44,201; John Sopko, Reg. No. 41,321, and Barry Jacobsen, Reg. No. 43,689, each of them of HOFFMANN & BARON, LLP, 1055 Parsippany Boulevard, Parsippany, New Jersey 07054.

PLEASE SEND CORRESPONDENCE TO:

Irving N. Felt, Esq.
HOFFMANN & BARON, LLP
6900 Jericho Turnpike
Syosset, NY 11791

PLEASE DIRECT TELEPHONE CALLS TO:

James F. Harrington, Esq.

(516) 822-3550

DECLARATION

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

SIGNATURE(S)


1-00 Full Name of Sole or First Inventor: Jean-Gérard Guillet

Country of Citizenship: France

Residence Address: 39, Rue Raphaél, Vanves, F-92170 France FLX

Post Office Address: Same as Above

Date: 28/2/2000

Inventor's signature 

2-00 Full Name of Second Joint Inventor: Gilles Guichard


Country of Citizenship: France

Residence Address: 6, Quai Matthis, Strasbourg, F-67000 France

LA LOUVIERE - 5, rue du Milieu - 67200 WOLFISHEIM FLX

Post Office Address: Same as Above

Date: 03/02/2000

Inventor's signature 


3-00 Full Name of Third Joint Inventor: Marina Ostankovitch

Country of Citizenship: France

Residence Address: 89, Rue Broca, Paris F-75013 France FLX

Post Office Address: Same as Above

Date: 02 22 00

Inventor's signature 


4) 00
Full Name of Fourth Joint Inventor: Francine Connan
Country of Citizenship: France
Residence Address: 21, Rue Du Progrès, Deuil-La-Barre, F-95170 France FLX
Post Office Address: Same as Above
Date: 28/2/2000 Inventor's signature: F. Connan

500
Full Name of Fifth Joint Inventor: Anne Quesnel
Country of Citizenship: France
Residence Address: 13A, Boulevard De Lyon, Strasbourg, F-67000, France FLX
8, rue de Spesbourg
Post Office Address: Same as Above
Date: 20/03/2000 Inventor's signature: A. Quesnel

600
Full Name of Sixth Joint Inventor: Jeannine Choppin
Country of Citizenship: France
Residence Address: 8, Rue Saint Fiacre, Reims, F-51100 France FLX
Post Office Address: Same as Above
Date: 28/2/2000 Inventor's signature: J. Choppin

700
Full Name of Seventh Joint Inventor: Jean-Paul Briand
Country of Citizenship: France
Residence Address: 11, rue Beethoven
22, Rue Des Balayeurs, Strasbourg, F-67000 France FLX
Post Office Address: Same as Above
Date: 3/3/2000 Inventor's signature: J. Briand

8-012

Full Name of Eighth Joint Inventor: Sylviane Muller
Country of Citizenship: France
Residence Address: 15, Avenue De La Forêt Noire, Strasbourg, F-67000, France ~~11, rue Beethoven~~ FX
Post Office Address: Same as Above
Date: March 3, 2000 Inventor's signature 

NOTE All above spaces identifying inventors must be completed or deleted before any inventor executes this application

103372_1.DOC